

Primitive Ideals of Noetherian Generalized Down-Up Algebras

Iwan Praton

Abstract

We classify the primitive ideals of noetherian generalized down-up algebras.

Keywords: down-up algebra; generalized down-up algebra; enveloping algebra; primitive ideal.

2000 Mathematics Subject Classification: 16D60; 17B35; 16D25; 16S36.

1 Introduction and preliminaries

1.1 Background

We can construct $U(\mathfrak{sl}_2)$, the enveloping algebra of $\mathfrak{sl}_2$, as the algebra generated by three elements h, x, y , subject to the relations

$$hx - xh = x, \quad yh - hy = y, \quad xy - yx = h.$$

(The first two relations are somewhat nontraditional.) There are various ways of generalizing this construction. One could increase the number of generators; this could lead to enveloping algebras of other Lie algebras. Another path would be to stay with three generators but modify the defining relations. We would then get an algebra with a reasonably small dimension, making the algebra computationally tractable.

One such example is Smith's algebras similar to $U(\mathfrak{sl}_2)$; see [S]. These algebras are generated by h, x, y subject to

$$hx - xh = x, \quad yh - hy = y, \quad xy - yx = \phi(h),$$

where ϕ is a polynomial. Thus h in this case is no longer the commutator of x and y , but a polynomial root of said commutator. The resulting algebras share many properties with $U(\mathfrak{sl}_2)$.

Another example, going in a different direction, are the down-up algebras of Benkart and Roby [BR]. These can be defined as the algebras generated by h, u, d subject to the relations

$$hu - ruh = \gamma u, \quad dh - rhd = \gamma d, \quad du - sud = h,$$

where r, s , and γ are constants. When $\gamma \neq 0$, these algebras have similar defining relations with $U(\mathfrak{sl}_2)$, except the commutators have been modified. Down-up algebras, especially when $rs\gamma \neq 0$, also share many properties with $U(\mathfrak{sl}_2)$.

Other generalizations of $U(\mathfrak{sl}_2)$ exist, some of which can be considered as a mixture of Smith's algebras and down-up algebras. For example, Rueda has studied algebras that are generated by h, x, y subject to

$$hx - xh = x, \quad yh - hy = y, \quad xy - sxy = \phi(h),$$

where s is a constant and ϕ is a polynomial [R]. For another example, consider the conformal $\mathfrak{sl}_2$ enveloping algebras of Le Bruyn [LB], which are generated by h, u, d subject to

$$hu - ruh = u, \quad dh - rhd = d, \quad du - sud = ah^2 + h,$$

where r, s, a are constants with $rs \neq 0$.

In 2004 Cassidy and Shelton introduced the ultimate mixture of Smith's algebras and down-up algebras [CS]. These algebras are generated by h, u, d subject to

$$hu - ruh = \gamma u, \quad dh - rhd = \gamma d, \quad du - sud = \phi(h),$$

where r, s, γ are constants and ϕ is a polynomial. It is these algebras that are the subject of this paper.

More specifically, we classify the primitive ideals of noetherian generalized down-up algebras, hence completing the project begun in [P2] and [P3]. As before, we try to provide a reasonably explicit list of generators for these primitive ideals. Most of the necessary techniques are straightforward generalizations of [P1]. In particular, most of the time we will be using explicit computation—part of the charm of studying down-up algebras is that elementary computations actually lead to useful results; it is not necessary to rely on heavy theoretical machinery.

1.2 Notations

As usual, we denote the complex numbers by $\mathbb{C}$; $\mathbb{C}^\times$ will denote the nonzero complex numbers. If $z \in \mathbb{C}$, it is convenient to define $o(z)$ to be the order of z in the multiplicative group $\mathbb{C}^\times$. Thus z is a primitive n th root of unity if and only if $o(z) = n$. If z is not a root of unity, we set $o(z) = \infty$. By convention, when we write $o(z) = n$ or m , we take n or m to be finite, i.e., z is a root of unity.

We also need to use an ordering on the degree of two-variable polynomials, so we describe here the ordering that we will use. The monomial $x^i y^j$ has degree (i, j) ; the degrees are ordered “alphabetically by last name”, i.e., $(i, j) > (i', j')$ iff $j > j'$ or $j = j'$ and $i > i'$. The degree of a polynomial is the highest degree of its constituent monomials.

We write $\langle a, b, \dots, z \rangle$ to denote the (two-sided) ideal generated by the elements $a, b, \dots, z$.

1.3 Definition of generalized down-up algebras

We now state a careful definition of our object of study. A generalized down-up algebra is an algebra over $\mathbb{C}$ parametrized by three complex numbers and a complex polynomial. Specifically, the algebra $L(\phi, r, s, \gamma)$, where $r, s, \gamma \in \mathbb{C}$ and $\phi \in \mathbb{C}[x]$, is the $\mathbb{C}$ -algebra generated by three generators u , d , and h , subject to the relations

$$hu - ruh = \gamma u, \quad dh - rhd = \gamma d, \quad du - sud = \phi(h).$$

(We follow the convention in [P3], which is slightly different from [CS].) We often just write L for $L(\phi, r, s, \gamma)$ when the parameter values are implicitly known. The algebra L is noetherian if and only if $rs \neq 0$. Primitive ideals in the non-noetherian case was described in [P3], so in this paper we always assume that $rs \neq 0$. This assumption implies that L is a domain [CS, Proposition 2.5].

1.4 Bases

In order to do computations, we need a basis for L . The standard basis consists of the monomials $\{u^i h^j d^k : i, j, k \geq 0\}$. Since we are assuming that L is a domain, we can arrange the us , ds , and hs in any order, i.e., the monomials $\{u^i d^k h^j\}$, $\{d^k h^j u^i\}$, $\{d^k u^i h^j\}$, $\{h^j u^i d^k\}$, and $\{h^j d^k u^i\}$ (where $i, j, k \geq 0$) are all bases of L [CS, Theorem 2.1].

1.5 Grading

There is a useful grading on L that results from declaring that u has degree $+1$, d has degree -1 , and h has degree 0 . (Thus the monomial $u^i h^j d^k$ has degree $i - k$.) Clearly L_0 , the elements of degree 0 , is itself an algebra. It turns out to be a polynomial algebra on the two variables h and ud [CS, Proposition 4.1]. If $i > 0$, then any element in L_i can be written (uniquely) as $u^i f(h, ud)$, where $f \in \mathbb{C}[x, y]$. Similarly, if $k < 0$, then any element in L_k can be written (uniquely) as $g(h, ud)d^k$, where $g \in \mathbb{C}[x, y]$.

As in the commutative case, we say that an element of L_i is *homogeneous of degree i* . Any $x \in L$ can be written as a sum of homogeneous elements: $x = \sum_{i \in \mathbb{Z}} x_i$, where $x_i \in L_i$ and only finitely many of the x_i s are nonzero. We define the length $\ell(x)$ of x to be the number of nonzero x_i s: $\ell(x) = \#\{i \in \mathbb{Z} : x_i \neq 0\}$. (Thus an element of length 1 is a nonzero homogeneous element.)

1.6 Isomorphisms

Different values of the parameters ϕ , r , s , and γ can lead to isomorphic algebras. For example, $L(\phi, r, s, \gamma)$ and $L(\psi, r, s, c\gamma)$ are isomorphic, where $c \neq 0$ and $\psi(x) = \phi(cx)$. (The isomorphism sends u to u' , d to d' , and h to ch' .) Thus we can assume that either $\gamma = 0$ or $\gamma = 1$ without loss of generality. Similarly, there is an isomorphism between $L(\phi, r, s, \gamma)$ and $L(c\phi, r, s, \gamma)$ via $u \mapsto cu'$, $d \mapsto d'$, and $h \mapsto h'$. Thus we can assume that the polynomial ϕ is either monic or zero. We will, however, continue to use γ and ϕ without additional assumptions, since assuming $\gamma = 1$ or ϕ monic does not significantly ease our workload. But there *is* an isomorphism between generalized down-up algebras that we will exploit heavily; see [CL, Proposition 1.7]

Lemma 1.1. *Suppose $r \neq 1$. Then $L(\phi, r, s, \gamma)$ is isomorphic to $L(\psi, r, s, 0)$, where $\psi(x) = \phi(x - \gamma/(r - 1))$.*

This lemma clarifies the role of γ : in most cases, γ is not necessary! The information carried by γ can be transferred into the polynomial ϕ . Note that this phenomenon is not visible in the original formulation of down-up algebras, since in that setting we do not have flexibility in the choice of ϕ .

We will thus treat the cases $r = 1$ and $r \neq 1$ differently. When $r = 1$, we will consider both $\gamma = 0$ and $\gamma \neq 0$. But when $r \neq 1$, we will assume that $\gamma = 0$; this cuts down the number of cases we have to consider by almost half.

1.7 Conformal algebras

Here is an example of the usefulness of Lemma 1.1. Recall that the algebra $L(\phi, r, s, \gamma)$ is called *conformal* if there exists a polynomial ψ such that $s\psi(x) - \psi(rx + \gamma) = \phi(x)$. Conformal algebras are nice because we can then define $H = ud + \psi(h)$; we shall see that having such an element is quite useful. For one, any polynomial in ud and h can be written as a polynomial in H and h ; the commutation relations involving H are more convenient than those involving ud . Specifically, it is straightforward to show that $Hu = suH$ and $dH = sHd$.

To determine exactly when an algebra is conformal is not a complete triviality, mostly due to the presence of γ , but Lemma 1.1 allows us to ignore γ most of the time, so we can determine quickly which of our algebras are conformal [CL, Lemma 1.6, Proposition 1.8].

Lemma 1.2. *Suppose $\phi(x) = \sum_{i=0}^n a_i x^i$. If $s \neq r^j$ for $0 \leq j \leq n$, then $L(\phi, r, s, 0)$ is conformal. If $s = r^j$ for some $0 \leq j \leq n$ and $a_j = 0$, then $L(\phi, r, s, 0)$ is also conformal. Otherwise $L(\phi, r, s, 0)$ is not conformal. Additionally, when $\gamma \neq 0$, $L(\phi, 1, s, \gamma)$ is conformal.*

Since conformal algebras behave somewhat differently than nonconformal ones, we will treat these two cases separately. It turns out that the nonconformal case, even when $\gamma = 0$, has a similar flavor to the situation when $\gamma \neq 0$.

1.8 Schur's Lemma

Finally, one of our main weapons is the following result, well-known to representation theorists as Dixmier's version of Schur's lemma [D, 2.6.5].

Lemma 1.3. *Let A be an $\mathbb{C}$ -algebra and M a simple A -module whose dimension is countable. If $\xi \in \text{hom}_{\mathbb{C}}(M, M)$ commutes with the action of A on M , then ξ acts as a scalar on M .*

In particular, the center of A acts as scalar operators on M . We will use this result so often that we will not mention it explicitly.

2 Weight modules and finite dimensional simple modules

In this section we describe weight modules; for our purposes, we are especially interested in universal weight modules and finite-dimensional modules

(which are instances of weight modules). When simple, these modules provide almost all of our primitive ideals. In some cases, we do have to consider modules that are not weight modules, but such cases are exceptional, and we will deal with them when they arise.

Finite dimensional simple modules have been classified in [CS, Section 4], so all we have to do here is figure out their annihilators.

2.1 Universal weight Modules

We first recall the definition of weight modules. If M is an L -module, then $v \in M$ is said to have weight $(\lambda, \beta) \in \mathbb{C}^2$ if $h \cdot v = \lambda v$ and $(ud) \cdot v = \beta v$. The *weight space* $M_{(\lambda, \beta)}$ is the linear space consisting of all elements of M with weight (λ, β) . The module M is a *weight module* if it is the (direct) sum of its weight spaces.

There is a nice relation between weight modules and the grading of L described in section 1.5. We need to recall the (invertible) operation on weights given by $\Phi : (\lambda, \beta) \mapsto (r\lambda + \gamma, s\beta + \phi(\lambda))$. (See [CS, section 4].) Then a simple calculation shows that if M is a weight module, then $L_i M_{(\lambda, \beta)} \subseteq M_{\Phi^i(\lambda, \beta)}$ ($i \in \mathbb{Z}$), i.e., elements of degree i transform vectors of weight (λ, β) into vectors whose weights is i steps away.

There exist universal weight modules, which we now describe. Let (λ, β) be an arbitrary weight, and define $(\lambda_i, \beta_i) = \Phi^i(\lambda, \beta)$ ($i \in \mathbb{Z}$) as above. Then the universal weight module $W(\lambda, \beta)$ is the module with basis $\{v_i : i \in \mathbb{Z}\}$ and

$$\begin{aligned} hv_i &= \lambda_i v_i, & i \in \mathbb{Z}; \\ uv_i &= v_{i+1}, & dv_{-i} = v_{-i-1}, & i \geq 0; \\ uv_{-i} &= \beta_{-i+1} v_{-i+1}, & dv_i &= \beta_i v_{i-1}, & i > 0. \end{aligned}$$

Any weight module is a quotient of a universal weight module. (It is possible to describe universal weight modules more succinctly as a tensor product, but since we want to do explicit calculations on these modules, it is better to have explicit formulas for the action of L .)

We will pay a lot of attention to whether $W(\lambda, \beta)$ is simple. A straightforward result along these lines is as follows. If the weights (λ_i, β_i) are all distinct, and $\beta_i \neq 0$ for all $i \in \mathbb{Z}$, then $W(\lambda, \beta)$ is simple.

2.2 Finite-dimensional modules

We now turn to finite-dimensional modules. All simple finite-dimensional modules are weight modules, and hence quotients of the universal weight

modules. We need, however, to have more detailed information. It turns out that there are two types of finite-dimensional simple modules: those with highest (and lowest) weights, and those that are cyclic.

The highest weight simple modules can be described as follows. Start with a weight $(\lambda, 0)$, and as before write (λ_i, β_i) for $\Phi^i(\lambda, 0)$. (Thus $\lambda_0 = \lambda$ and $\beta_0 = 0$ in this notation.) Suppose the weights (λ_i, β_i) are all distinct, $\beta_{n+1} = 0$, and $\beta_i \neq 0$ for $1 \leq i \leq n$. Then there is a simple module of dimension $n + 1$, say with basis $\{v_0, v_1, \dots, v_n\}$. Each v_i is a vector with weight (λ_i, β_i) . The action of L on this module is

$$\begin{aligned} hv_i &= \lambda_i v_i, & 0 \leq i \leq n; \\ dv_0 &= 0, & dv_i = \beta_i v_{i-1}, & 1 \leq i \leq n; \\ uv_n &= 0, & uv_i = v_{i+1}, & 0 \leq i \leq n-1. \end{aligned}$$

We denote this simple module by $F_{\text{hw}}(\lambda)$, similar to the notation in [CS].

There are two types of cyclic simple modules, denoted by $F_c(\zeta, \rho)$ and $\overline{F}_c(\zeta, \rho)$ in [CS, Definition 4.6]. We describe $F_c(\zeta, \rho)$ first. Start with a weight (λ_0, β_0) , and suppose its orbit under Φ is finite; say the orbit has m distinct values (λ_i, β_i) , $0 \leq i \leq m-1$. Assume that $(\lambda_i, \beta_i) \neq (0, 0)$ for any i . Denote this set of weights by ζ . Let ρ be a nonzero complex number. Then there is a simple module $F_c(\zeta, \rho)$ of dimension m with basis, say, $\{v_0, v_1, \dots, v_{m-1}\}$, where the basis vectors are indexed by the cyclic group $\mathbb{Z}/m\mathbb{Z}$. Thus, for example, v_{m+1} is the same as v_1 . Each v_i is vector of weight (λ_i, β_i) . Then action of L on $F_c(\zeta, \rho)$ is

$$hv_i = \lambda_i v_i, \quad uv_i = \rho v_{i+1}, \quad dv_i = \beta_i v_{i-1}.$$

The modules $\overline{F}_c(\zeta, \rho)$ are similar. The parameters and the weights are the same; the only difference with $F_c(\zeta, \rho)$ is in the action of L :

$$hv_i = \lambda_i v_i, \quad uv_i = \beta_{i+1} v_{i+1}, \quad dv_i = \rho v_{i-1}.$$

There is usually a lot of overlap between $F_c(\zeta, \rho)$ and $\overline{F}_c(\zeta, \rho')$ as ρ and ρ' vary over $\mathbb{C}^\times$, but we need to consider both types of cyclic modules since $F_c(\zeta, \rho)$ and $\overline{F}_c(\zeta, \rho')$ are nonisomorphic when $\prod_{i \in \mathbb{Z}/m\mathbb{Z}} \beta_i = 0$.

2.3 Annihilators of finite-dimensional simple modules

We now figure out the annihilators of these simple modules. First we look at $F_{\text{hw}}(\lambda)$. In this case, let $J_\lambda = \{f \in L_0 = \mathbb{C}[h, ud] : f(\lambda_i, \beta_i) = 0, 0 \leq i \leq n\}$. Then J_λ is a classical polynomial ideal; it is finitely generated, and for specific values of (λ_i, β_i) we can figure out a list of its generators.

Proposition 2.1. *The annihilator of $F_{hw}(\lambda)$ is $\langle u^{n+1}, d^{n+1}, J_\lambda \rangle$.*

Proof. Write I for the ideal $\langle u^{n+1}, d^{n+1}, J_\lambda \rangle$. It is straightforward to verify that I annihilates $F_{hw}(\lambda)$; we need to show the reverse inclusion $\text{Ann } F_{hw}(\lambda) \subseteq I$.

If $n = 0$, then $F_{hw}(\lambda)$ is one dimensional; u and d act as the zero operator while h acts as the scalar λ . The ideal J_λ is generated by ud and $h - \lambda$, so in this case, $I = \langle u, d, h - \lambda \rangle$. It is clear that I is the annihilator of $F_{hw}(\lambda)$. Thus we assume that $n \geq 1$ from now on.

Now suppose y is an element of L that annihilates $F_{hw}(\lambda)$. Since $L_i v_j \subseteq \mathbb{C}v_{j+i}$, it does no harm to assume that y is homogeneous, say of degree $-k < 0$. (If y is homogeneous of positive degree, the proof is similar.)

We'll utilize the element $x = u + d^n$. Note that $xv_i = v_{i+1}$ for $0 \leq i \leq n-1$, but $xv_n = \beta_1 \cdots \beta_n v_0$; thus $x^n v_i = \eta v_i$, where $\eta = \prod_{i=1}^n \beta_i$ is nonzero. So x^n acts a nonzero scalar on $F_{hw}(\lambda)$.

Note also that $x^2 = u^2 + ud^n + d^n u + d^{2n} \in u^2 + L_{1-n} + I$; by induction we can show that $x^i \in u^i + L_{i-1-n} + I$ for $1 \leq i \leq n$.

Recall that $y \in d^k L_0$. Thus $yx^k \in (u^k + L_{k-1-n} + I)(d^k L_0) \in L_0 + I$. In other words, $yx^k \in f(h, ud) + I$, where f is a polynomial in two variables. But yx^k annihilates everything in $F_{hw}(\lambda)$, so $f(h, ud)$ also annihilates everything, and hence $f(h, ud)$ must be in J_λ . So yx^k is actually in I .

Thus modulo I , we have $0 \equiv yx^k \equiv yx^k x^{n-k} \equiv yx^n \equiv \eta y$. Since $\eta \neq 0$, we conclude that $y \equiv 0$, which is what we want to show. $\square$

We now take a look at the annihilators of $F_c(\zeta, \rho)$ and $\overline{F}_c(\zeta, \rho)$. We again utilize the ideal $J_\zeta = \{f(h, ud) : f(\lambda_i, \beta_i) = 0, i \in \mathbb{Z}/m\mathbb{Z}\}$ in L_0 . Write $\eta = \prod_{i=0}^{m-1} \beta_i$.

Proposition 2.2. *The annihilator of $F_c(\zeta, \rho)$ is $\langle u^m - \rho^m, d^m - \eta, J_\zeta \rangle$, and the annihilator of $\overline{F}_c(\zeta, \rho)$ is $\langle d^m - \rho^m, u^m - \eta, J_\zeta \rangle$.*

Proof. The proof is similar to the previous proof. As before, it is clear that I annihilates $F_c(\zeta, \rho)$. We want to show that $\text{Ann } F_c(\zeta, \rho) \subseteq I$.

Write $L'_k = \sum_{i \equiv k \pmod m} L_i$. Then $L'_k v_i \subseteq \mathbb{C}v_{i+k}$. Suppose now that y annihilates $F_c(\zeta, \rho)$. It does no harm to assume that $y \in L'_k$ for some $k \in \mathbb{Z}$. Since $u^{am+k} \in \rho^{am} u^k + I$ and $d^{bm+(m-k)} \in \eta^b d^{m-k}$, we can even assume that $y \in u^k L_0 + d^{m-k} L_0$ where $0 \leq k \leq m-1$.

Now yu^{m-k} also annihilates $F_c(\zeta, \rho)$, and $yu^{m-k} \in L_m + L_0 \subseteq L_0 + I$. Therefore we can write $yu^{m-k} = f(h, ud) + I$ where f is a polynomial of two variables. Since yu^{m-k} annihilates everything, we conclude that $f \in J_\zeta$, i.e., $yu^{m-k} \in I$.

Therefore $yu^{m-k}u^k$ is also in I , and so modulo I , we have $0 \equiv yu^m \equiv y\rho^m$. Because $\rho \neq 0$, we conclude that $y \equiv 0$, as required.

The proof for $\overline{F}_c(\zeta, \rho)$ is similar. $\square$

2.4 Detecting finite-dimensionality

It is also useful to be able to tell whether a simple module is finite-dimensional from partial information about its annihilator.

Lemma 2.3. *Let M be a simple module such that (i) either d^m or u^m ($m \geq 1$) acts as a scalar on M ; (ii) M contains a weight vector. Then M is finite-dimensional.*

Proof. We suppose that d^m acts as the scalar ρ^m . (The proof where u^m acts as a scalar is similar.) We first consider the case where $\rho = 0$.

In this case the first step is to show that M contains an element with weight $(\lambda, 0)$. Recall that if v is a weight vector, then $d^i v$ is also a weight vector, for all $i \geq 0$. Since $d^m v = 0$, there must be an $j \geq 0$ such that $d^j v \neq 0$ and $d^{j+1} v = 0$. Then $w_0 = d^j v$ is an element of weight $(\lambda, 0)$ for some $\lambda \in \mathbb{C}$.

Now define $w_i = u^i w_0$, $i \geq 0$. It is straightforward to check that $\text{span}\{w_i : i \geq 0\}$ is stable under u , d , and h ; since M is simple, we have $M = \text{span}\{w_i : i \geq 0\}$.

Note that $dw_i = \beta_i w_{i-1}$ for some $\beta_i \in \mathbb{C}$. Thus we have $0 = d^m w_m = \beta_m \beta_{m-1} \cdots \beta_1 w_0$. Since $w_0 \neq 0$, we have $\beta_k = 0$ for some $1 \leq k \leq m$. Then $M_0 = \text{span}\{w_{k+i} : i \geq 0\}$ is a submodule of M , and hence must be 0 or M . If $M_0 = 0$, then M is spanned by $\{w_0, \dots, w_{k-1}\}$ and so is finite-dimensional. If $M_0 = M$, then in particular w_0 is a linear combination of the elements w_{k+i} , $i \geq 0$. Say $w_0 = \sum_{i=1}^t c_i w_{n_i}$, where $c_i \neq 0$ and we arrange the indices so that $n_i > n_j$ whenever $i > j$. Then w_{n_t} is a linear combination of w_0 and the elements w_{n_i} , $i < t$. Applying the operator u , we see that any w_j , with $j \geq n_t$, is a linear combination of elements w_i , with $i < n_t$. Thus M is spanned by $\{w_i : i < n_t\}$, and hence finite-dimensional.

We now consider the case where $\rho \neq 0$. As before, suppose $v \in M$ is a weight vector. Let $v_i = u^i v$ for $0 \leq i \leq m-1$. Each v_i is still a weight vector; in particular, $h v_i \in \mathbb{C} v_i$ for all values of i . Since v_0 is an eigenvector of ud , we check easily that $dv_i \in \mathbb{C} v_{i-1}$ for $1 \leq i \leq m-1$. Since u^m acts as ρ^m on M , we get that $uv_{m-1} = \rho^m v_0$. Thus $dv_0 = \rho^{-m} duv_{m-1} = \rho^{-m} [sud + \phi(h)]v_{m-1} \in \mathbb{C} v_{m-1}$. Therefore the span of $\{v_i : 0 \leq i \leq m-1\}$ is stable under L , and hence must be all of M . Thus M is finite-dimensional. $\square$

Corollary 2.4. *Let M be a simple module, and suppose for some $m \geq 1$, we have $d^m \in \text{Ann } M$, $d^{m-1} \notin \text{Ann } M$, and $f(h)d^{m-1}M = 0$ for some nonconstant polynomial f . Then M is finite-dimensional. (The same conclusion holds if d is replaced by u .)*

In particular, if $d^m u - s^m u d^m \neq 0$, then M is finite-dimensional. (Similarly, if $u^m \in \text{Ann } M$, $u^{m-1} \notin \text{Ann } M$, and $du^m - s^m u^m d \neq 0$, then M is finite-dimensional.)

Proof. Suppose f has degree $n \geq 1$. Pick a nonzero element $v \in d^{m-1}M$, and let $W = \text{span}\{v, hv, h^2v, \dots, h^{n-1}v\}$. Then W is stable under h , and since W is finite-dimensional, h has an eigenvector in W , say w . Note that $dhv = (rh + \gamma)dv = 0$; similarly, $dh^i v = 0$ for all $i \geq 0$. Thus $dw = 0$, and so w is a weight vector. We can now apply the lemma.

In the particular case, we show easily by induction that, for $k \geq 1$,

$$d^k u - s^k u d^k = f_k(h) d^{k-1}$$

for some polynomial f_k . The hypothesis implies that $f_m(h) \neq 0$. Note that $f_m(h)$ cannot be a constant since that would imply $f_m(h)d^{m-1}M = 0$, i.e., $d^{m-1}M = 0$. Thus $f_m(h)$ is a nonconstant polynomial, and the conclusion follows. $\square$

It is also useful occasionally to detect when a simple module is one-dimensional.

Lemma 2.5. *Let M be a simple module. Suppose ud and h act as scalars on M . Then M is one-dimensional.*

Proof. Say ud acts as the scalar β and h acts the scalar λ . Pick a nonzero $w_0 \in M$ and for $i \geq 0$ define $w_i = u^i w_0$ and $w_{-i} = d^i w_0$. For all $i \in \mathbb{Z}$, we clearly have $uw_i \in \mathbb{C}v_{i+1}$; also, since $du = sud + \phi(h)$, we get that $dv_i \in \mathbb{C}v_{i-1}$ for $i \in \mathbb{Z}$. Thus $\text{span}\{w_i : i \in \mathbb{Z}\}$ is a submodule of M and hence must be all of M .

If $i < 0$, we have $uw_i = udw_{i+1} = \beta w_{i+1}$, so $duw_i = \beta w_i$. If $i \geq 0$, then $uduw_i = \beta uw_i = \beta w_{i+1}$; but we also know that $duw_i = \alpha w_i$ for some $\alpha \in \mathbb{C}$, so $uduw_i = \alpha w_{i+1}$. Thus $\alpha = \beta$ and we conclude that $duw_i = \beta w_i$.

Therefore ud and du act as the same scalar on M . Thus the operators u , d , and h all commute with each other, and hence M is one-dimensional. $\square$

3 When $r = 1$ and $\gamma \neq 0$

In this section we tackle the case where $\gamma \neq 0$. Recall from Lemma 1.1 that in this case it suffices to assume that $r = 1$. We then have $hu = u(h + \gamma)$ and

$hd = d(h - \gamma)$. It follows easily that $f(h)u = uf(h + \gamma)$ and $f(h)d = df(h - \gamma)$ where f is any polynomial; also, $hu^i = u^i(h + i\gamma)$ and $hd^i = (h - i\gamma)d^i$ for $i \geq 0$. We can summarize succinctly by saying that $f(h)x_i = x_if(h + i\gamma)$, where $x_i \in L_i$.

3.1 Homogeneous elements in ideals

We start with a result about homogeneous elements.

Lemma 3.1. *Let I be an ideal of L . Suppose $x \in I$ and $x = \sum_{i \in \mathbb{Z}} x_i$ (where $x_i \in L_i$) is the homogeneous decomposition of x . Then $x_i \in I$ for all $i \in \mathbb{Z}$.*

Proof. We use induction on the length of x . (Recall that the length of x is the number of nonzero x_i s.) The lemma is certainly true for elements of length 1. Assume it is true for elements of length $n - 1$, and let $x = \sum_{i \in \mathbb{Z}} x_i$, ($x_i \in L_i$) be an element of length n in I . Thus exactly n of the x_i s is nonzero. Pick an integer m so that $x_m \neq 0$. Now

$$hx = \sum_{i \in \mathbb{Z}} hx_i = \sum_{i \in \mathbb{Z}} x_i(h + i\gamma),$$

so

$$xh + m\gamma x - hx = \sum_{i \in \mathbb{Z}} (m - i)\gamma x_i.$$

This is an element in I of length $n - 1$. We apply the induction hypothesis and conclude that $(m - i)\gamma x_i \in I$ for $i \in \mathbb{Z}$, and thus $x_i \in I$ for $i \neq m$. Therefore $y = \sum_{i \neq m} x_i$ is an element of I , hence so is $x_m = x - y$. $\square$

Recall from Lemma 1.2 that the algebra $L(\phi, 1, s, \gamma)$ is conformal. Thus there exists a polynomial $\psi \in \mathbb{C}[x]$ such that $s\psi(x) - \psi(x + \gamma) = \phi(x)$. Recall also that we define H as the element $ud + \psi(h)$; then $Hu = suH$ and $dH = sHd$. We conclude that $f(h, H)u = uf(h + \gamma, sH)$ and $f(h, H)d = df(h - \gamma, s^{-1}H)$ for any polynomial $f \in \mathbb{C}[x, y]$.

Note that if M is an L -module, then HM is stable under u and d (and certainly under h), so HM is a submodule. Thus if M is simple, then either $HM = 0$ or $HM = M$. We treat these two cases separately.

3.2 The case $HM = 0$

Suppose first that $HM = 0$. Of course, $\text{Ann } M$ then contains $\langle H \rangle$, but if ψ is the zero polynomial, we can say more. In this case, $du = sud$, so uM and dM are submodules of M , and hence either $uM = 0$ or $uM = M$; similarly,

either $dM = 0$ or $dM = M$. It cannot be the case that both $uM = M$ and $dM = M$, for then $HM = udM = M$, contradicting $HM = 0$. Thus when ψ is the zero polynomial (and $HM = 0$), $\text{Ann } M$ must contain either u or d .

Lemma 3.2. *Suppose M is simple and $HM = 0$. If $\text{Ann } M \supsetneq \langle H \rangle$, then $\text{Ann } M$ contains a power of u or a power of d . Furthermore, if ψ is the zero polynomial and $\text{Ann } M \supsetneq \langle d \rangle$, then $\text{Ann } M$ contains a power of u , while if $\text{Ann } M \supsetneq \langle u \rangle$, then $\text{Ann } M$ contains a power of d .*

Proof. When ψ is not the zero polynomial, let $I = \langle H \rangle$, but when ψ is the zero polynomial, let I denote either $\langle u \rangle$ or $\langle d \rangle$. Suppose that $\text{Ann } M$ strictly contains I . Choose an element $x \in \text{Ann } M$ that is not in I ; by Lemma 3.1 we can assume that x is homogeneous, say of degree k . If $I = \langle u \rangle$, then $k \leq 0$; if $I = \langle d \rangle$, then $k \geq 0$; if $I = \langle H \rangle$, then there is no restriction on k . To avoid repetitions, we assume $k \geq 0$ here (and thus $I = \langle d \rangle$ if $\psi = 0$); the other possibilities are treated similarly. Then we can write $x = u^k g(h, H)$, but since $x \notin I$, we can assume that $x = u^k f(h)$ for some polynomial f .

Among all nonzero elements of the form $u^k f(h)$, pick one where the degree of f is as small as possible. Then

$$xu - ux = u^{k+1}(f(h + \gamma) - f(h)).$$

The polynomial $f(h + \gamma) - f(h)$ has smaller degree than $f(h)$, hence it must be the zero polynomial by choice of x . Thus we have $f(h + \gamma) = f(h)$, and this implies that f is the constant polynomial. Thus $\text{Ann } M$ contains u^k , where $k \geq 1$. $\square$

We can now derive some consequences about M .

Lemma 3.3. *Suppose M is simple and $HM = 0$. If ψ is not the zero polynomial and $\text{Ann } M \supsetneq \langle H \rangle$, then M is finite-dimensional. If ψ is the zero polynomial and $\text{Ann } M \supsetneq \langle u \rangle$ or $\text{Ann } M \supsetneq \langle d \rangle$, then M is one-dimensional.*

Proof. Suppose first that $\psi = 0$ and $\text{Ann } M \supsetneq \langle d \rangle$. Then the previous lemma implies that $u^k \in \text{Ann } M$ for some $k \geq 1$; this implies that $u \in \text{Ann } M$ (since either $uM = 0$ or $uM = M$). Hence $\text{Ann } M$ contains both u and d , and therefore all three operators u , d , and h commute, so they all act as scalars. Thus M is one-dimensional. Similarly, if $\psi = 0$ and $\text{Ann } M \supsetneq \langle u \rangle$, then M is one-dimensional.

On the other hand, suppose now that $\psi \neq 0$. By the previous lemma $\text{Ann } M$ contains u^k or d^k ; we'll say u^k for definiteness. We first show that

ψ cannot be a constant. If ψ were the constant C , then $ud = H - C$ would act as $-C$ on M , and thus $u^k d^k$ would act as the nonzero constant $(-C)^k$ on M , contradicting $u^k M = 0$. Thus ψ must be a nonconstant polynomial. Let u^m be the smallest power of u that lies in $\text{Ann } M$. In this case,

$$du^m - s^m u^m d = [s^m \psi(h - (m-1)\gamma) - \psi(h + \gamma)]u^{m-1}.$$

The polynomial $s^m \psi(h - (m-1)\gamma) - \psi(h + \gamma)$ is nonzero, since $s^m \psi(h - (m-1)\gamma) = \psi(h + \gamma)$ implies that ψ has infinitely many roots. (If α is a root of ψ , then so are $\alpha + m\gamma$, $\alpha + 2m\gamma$, and so on.) Thus we can use Corollary 2.4 to conclude that M is finite-dimensional. $\square$

3.3 The case $HM = M$

When $HM = M$, the situation is more complicated, especially when $o(s) < \infty$. If $o(s) = n$, then $H^n u = u H^n$ and $d H^n = H^n d$, so H^n is a central element in L . Thus H^n acts as a scalar c^n for some constant c . (We write the constant as an n th power for balance with H^n .) Thus any primitive ideal must contain $H^n - c^n$; since $HM = M$, we have $c \neq 0$. There is a special case, however, that we need to look at more closely. It is complicated enough that we bestow upon it a separate lemma.

Lemma 3.4. *Suppose $o(s) = n$, ψ is a constant polynomial $C \neq 0$, and H^n acts as the scalar C^n on a simple module M . Then $\text{Ann } M$ must contain either u^n or d^n .*

Proof. We have $\phi(h) = (s-1)C$, so $du = sud + (s-1)C$. By induction we get that $d^i u = s^i u d^i + (s^i - 1)C$ for $i \geq 1$. In particular, $d^n u = u d^n$. Thus $u d^n M = d^n u M$; also, $h d^n M = d^n (h - n\gamma) M$. Therefore $d^n M$ is submodule of M , hence either $d^n M = 0$ or $d^n M = M$. Similarly, either $u^n M = 0$ or $u^n M = M$. (Note that this result holds for any simple module M , not just those where H^n acts as C^n .)

Recall that $ud = H - C$. By induction we establish that $u^k d^k = \prod_{i=0}^{k-1} (s^{-i} H - C) = (-1)^k \prod_{i=0}^{k-1} (C - s^{-i} H)$. Thus $u^n d^n = (-1)^n \prod_{i=0}^{n-1} (C - s^{-i} H) = (-1)^n (C^n - H^n)$. Since H^n acts as C^n on M , we conclude that $u^n d^n$ acts as the zero operator on M . Therefore either $u^n M = 0$ or $d^n M = 0$, i.e., $\text{Ann } M$ contains either u^n or d^n . $\square$

We remark here that in the situation of the lemma, any ideal that contains u^n or d^n must also contain $H^n - C^n$, since $H^n - C^n = (-1)^{n+1} u^n d^n$.

Summarizing so far, we have the following information when $HM = M$: if $o(s) = n$, then $\text{Ann } M$ must contain $H^n - c^n$ for some $c \neq 0$; furthermore,

if $\psi = C \neq 0$ and $c^n = C^n$, then $\text{Ann } M$ actually contains u^n or d^n . In other cases (i.e., if s is not a root of unity), we have no information about $\text{Ann } M$.

We now look at primitive ideals that are possibly larger than the minimal ones. We start with a result about polynomials of two variables: it is the two-variable analogue of the fact that if $f(x+\gamma) = f(x)$, then f is a constant polynomial. (We used this fact in the proof of Lemma 3.3.)

Lemma 3.5. *Suppose $g(x, y) = \sum_{i=0}^m g_i(x)y^i$ with $m < o(s)$, and $g(x + \gamma, sy) = s^n g(x, y)$ for some $0 \leq n \leq m$. Then $g(x, y) = cy^n$ for some constant $c \in \mathbb{C}$.*

Proof. We have $\sum_{i=0}^m g_i(x + \gamma)s^i y^i = \sum_{i=0}^m s^n g_i(x)y^i$. Comparing the coefficients of y^i , we see that $s^i g_i(x + \gamma) = s^n g_i(x)$. Now if g_i has a root a , then $a + \gamma, a + 2\gamma, \dots$, are also roots. Thus g_i would have infinitely many roots, and hence must be the zero polynomial. If g_i does not have any roots, then it is a constant c , but then we would have $s^i c = s^n c$, so $c = 0$ unless $i = n$. We conclude that $g_i(x) = 0$ if $i \neq n$, but $g_n(x) = c$. Therefore $g(x, y) = cy^n$, as required. $\square$

Lemma 3.6. *Let M be a simple module with $HM = M$. If s is not a root of unity, and $\text{Ann } M \neq \{0\}$, then $\text{Ann } M$ contains a power of u or a power of d . If $o(s) = n$, and $\text{Ann } M$ strictly contains $\langle H^n - c^n \rangle$ for $c \neq 0$, then $\text{Ann } M$ also contains a power of u or a power of d .*

Additionally, suppose $\psi(h) = C \neq 0$ and $\text{Ann } M \supsetneq \langle d^n \rangle$. Then $u^n \in \text{Ann } M$. Similarly, if $\text{Ann } M \supsetneq \langle u^n \rangle$, then $d^n \in \text{Ann } M$.

Proof. If $o(s) = \infty$, let $I = \{0\}$; if $o(s) = n$, let $I = \langle H^n - c^n \rangle$, where $0 \neq c \in \mathbb{C}$. Suppose that $\text{Ann } M \supsetneq I$. We proceed as in the proof of Lemma 3.3. Choose an element $x \in \text{Ann } M$ that is not in I ; by Lemma 3.1 we can assume that x is homogeneous, say of degree k . For definiteness we take $k \geq 0$; the case $k < 0$ is similar. Thus $\text{Ann } M$ contains an element of the form $x = u^k g(h, H)$. If $o(s) = n$, i.e., $I = \langle H^n - c^n \rangle$, we can assume that the highest power of H that appears in g is at most $n - 1$. Among all nonzero elements of this form, choose one where the degree of g is minimal. Write $g(h, H) = \sum_{i=0}^m f_i(h)H^i$; here $m < o(s)$. Then $xu = u^k g(h, H)u = u^{k+1}g(h + \gamma, sH)$, so

$$xu - s^m ux = u^{k+1}[g(h + \gamma, sH) - s^m g(h, H)] = u^{k+1} \sum_{i=0}^m [s^i f_i(h + \gamma) - s^m f_i(h)] H^i.$$

The polynomial $g'(h, H) = \sum_{i=0}^m [s^i f_i(h + \gamma) - s^m f_i(h)] H^i$ has smaller degree than $g(h, H)$ (if $g(h, H)$ has degree (ℓ, m) , then $g'(h, H)$ has degree at most

$(\ell - 1, m)$), so by choice of $g(h, H)$, we must have $g'(h, H) = 0$. Hence $g(h + \gamma, sH) = s^m g(h, H)$, and by Lemma 3.5 we conclude that $g(h, H)$ is a nonzero multiple of H^m . Thus $\text{Ann } M$ contains $u^k H^m$. Since $HM = M$, we conclude that $\text{Ann } M$ contains u^k .

Now suppose $o(s) = n$, $\psi(h) = C \neq 0$, H^n acts as the scalar C^n on M , and $\text{Ann } M \supsetneq \langle d^n \rangle$. Note that $\langle d^n \rangle$ contains $d^{n-j} d^j u^j = d^{n-j} g_j(H)$, where $g_j(H) = \prod_{i=1}^j (s^i H - C)$ is a polynomial of degree j .

As before, we use Lemma 3.1 to conclude that $\text{Ann } M$ contains a homogeneous element $x \notin \langle d^n \rangle$ of degree k . Clearly $k > -n$. If $k \geq 0$, then we proceed as above, concluding that $\text{Ann } M$ contains a power of u . Suppose however that $k < 0$. Set $k = j - n$. Then we can write $x = d^{n-j} f(h, H)$; since $d^{n-j} g_j(H) \in I$, we can assume that the degree of H in $f(h, H)$ is at most $j - 1$. Therefore the element $g(h, H) = u^{n-j} x = u^{n-j} d^{n-j} f(h, H)$ is a nonzero element in $\text{Ann } M$; the degree of H in this element is at most $n - 1$. We can write $g(h, H) = \sum_{i=0}^m f_i(h) H^i$ where $m < n$. We now proceed as above, concluding that $\text{Ann } M$ contains a power of u . Since either $u^n M = 0$ or $u^n M = M$, we conclude that $u^n \in \text{Ann } M$.

Similarly, if $\text{Ann } M \supsetneq \langle u^n \rangle$, then $\text{Ann } M$ also contains d^n . $\square$

Lemma 3.7. *Let M be a simple module with $HM = M$.*

If $o(s) = \infty$ and $\text{Ann } M \neq \{0\}$, then M is finite-dimensional.

If $o(s) = n$, ψ is not a constant polynomial, and $\text{Ann } M \supsetneq \langle H^n - c^n \rangle$ ($c \in \mathbb{C}^\times$), then M is finite-dimensional.

If $o(s) = n$, $\psi(h) = C$ is constant, and $\text{Ann } M \supsetneq \langle H^n - c^n \rangle$ ($c \in \mathbb{C}^\times$), then $c^n = C^n$. (Thus $\text{Ann } M$ must contain u^n or d^n .)

If $o(s) = n$, $\psi(h) = C$ is constant, and $\text{Ann } M \supsetneq \langle u^n \rangle$ or $\text{Ann } M \supsetneq \langle d^n \rangle$, then M is finite-dimensional.

Proof. In all the cases listed, the preceding lemma implies that $\text{Ann } M$ contains a power of u or a power of d . For definiteness let's say that $u^k \in \text{Ann } M$.

Suppose now that $o(s) = \infty$ and $\text{Ann } M \neq \{0\}$. We first show that ψ cannot be a constant polynomial.

If ψ is the zero polynomial, then $H = ud$ and $u^k d^k = \prod_{i=0}^{k-1} s^{-i} H$ is in $\text{Ann } M$, i.e., H^k is in $\text{Ann } M$. This contradicts $HM = M$.

If ψ is a nonzero constant C , we have to work harder. Let v_0 be a nonzero element in $\ker u$ (such an element exists since $u^k M = 0$). Let $v_i = d^i v_0$. We check easily that for $i \geq 1$, $uv_i = (s^{-i} - 1)Cv_{i-1}$. In particular, $u^k v_k = cv_0$, where $c = C^k \prod_{i=1}^k (s^{-i} - 1) \neq 0$. This contradicts $u^k \in \text{Ann } M$.

Thus ψ is nonconstant. We now use Corrolary 2.4 to conclude that M is finite-dimensional.

Similarly, if $o(s) = n$, ψ is nonconstant and $\text{Ann } M \supsetneq \langle H^n - c^n \rangle$, we use Corollary 2.4 to conclude that M is finite-dimensional.

So it remains to consider the case where $o(s) = n$ and $\psi(h) = C$ is a constant polynomial. Now $u^k M = 0$ implies that $u^n \in \text{Ann } M$, so $u^n d^n = \pm(H^n - C^n)$ is also in $\text{Ann } M$. Thus the scalar $c^n - C^n$ is in $\text{Ann } M$, hence $c^n = C^n$, as required.

Suppose now that $\psi(h) = C$, $\text{Ann } M$ strictly contains $\langle u^n \rangle$ or $\langle d^n \rangle$. By the lemma above, $\text{Ann } M$ contains both u^n and d^n . We now show that if $u^n, d^n \in \text{Ann } M$, then M is finite-dimensional. First, choose a nonzero element v_0 such that $dv_0 = 0$ (such an element clearly exists). Define $v_i = u^i v_0$; then $v_j = 0$ for $j \geq n$ since $u^n \in \text{Ann } M$. Note that $Hv_0 = (ud + C)v_0 = Cv_0$. It follows that $Hv_i = Hu^i v_0 = s^i u^i H v_0 = s^i C v_i$ for $0 \leq i \leq n-1$. Since h commutes with H , we have $Hf_i(h)v_i = s^i f_i(h)v_i$ for any polynomial $f_i(h)$. Thus each $f_i(h)v_i$ belong to a different eigenspace of H , $0 \leq i \leq n-1$.

Now M is simple, so there exists an element $x \in L$ such that $xhv_0 = v_0$. We can rewrite this equation as $h \sum_{i=0}^{n-1} f_i(h)u^i v_0 = v_0$ for some polynomials f_i . Thus $h \sum_{i=0}^{n-1} f_i(h)v_i = v_0$. For $i \neq 0$, the H -eigenvalue of $f_i(h)v_i$ is different from the H -eigenvalue of v_0 , so we must have $f_i(h)v_i = 0$. Thus $hf_0(h)v_0 = v_0$, or $(hf_0(h) - 1)v_0 = 0$. In other words, we have found a nontrivial polynomial in h that annihilates v_0 . By Corollary 2.4, this implies that M is finite-dimensional. $\square$

3.4 Proving primitivity

We now show that the ideals $\{0\}$, $\langle H \rangle$, $\langle H^n - c^n \rangle$, $\langle u \rangle$, $\langle d \rangle$, $\langle u^n \rangle$, and $\langle d^n \rangle$ are all primitive, under the right circumstances.

Lemma 3.8. *The following are primitive ideals:*

1. $\langle H \rangle$, if ψ is not identically zero;
2. $\{0\}$, if $o(s) = \infty$;
3. $\langle H^n - c^n \rangle$ ($c \neq 0$), if $o(s) = n$ and ψ is not a constant polynomial;
4. $\langle H^n - c^n \rangle$ ($c \neq 0$), if $o(s) = n$ and ψ is a constant $C \neq c$;
5. $\langle u \rangle$ and $\langle d \rangle$, if $\psi(h) = 0$;
6. $\langle u^n \rangle$ and $\langle d^n \rangle$, if $o(s) = n$ and ψ is a nonzero constant C .

Proof. We make use of the universal weight module $W(\lambda, \beta)$ from section 2.1. In the present case ($r = 1, \gamma \neq 0$), we have $\lambda_i = \lambda + i\gamma$ and $\beta_i = s^i \beta + s^i \psi(\lambda) -$

$\psi(\lambda + i\gamma)$. Note that the λ_i s are always all distinct, so if $\beta_i \neq 0$ for all $i \in \mathbb{Z}$, then $W(\lambda, \beta)$ is simple.

We need to know how $H = ud + \psi(h)$ acts on $W(\lambda, \beta)$. A quick calculation shows that for $i \in \mathbb{Z}$, $Hv_i = s^i c$, where $c = \beta + \psi(\lambda)$. We then have $\beta_i = s^i c - \psi(\lambda + i\gamma)$. We usually set a value for c first, then find λ and β to suit our purpose.

Suppose first that ψ is a nonconstant polynomial. Then no matter what the value of c is, we can find λ and β such that $\beta + \psi(\lambda) = c$ and $\beta_i \neq 0$. (The details: For a given $i \in \mathbb{Z}$, the polynomial equation $\psi(x) = s^i c$ has only finitely many solutions, so there are only countably many values of λ where $\psi(\lambda + i\gamma) = s^i c$ for some $i \in \mathbb{Z}$. Thus we can certainly choose a value of λ so that $\psi(\lambda + i\gamma) \neq s^i c$ for *any* $i \in \mathbb{Z}$. We can then choose β to satisfy $\beta + \psi(\lambda) = c$.) Thus if ψ is a nonconstant polynomial, we can construct a simple $W(\lambda, \beta)$ for any value of c .

When $c = 0$, we get a simple module M with $HM = 0$. Additionally, note that even if ψ is a constant $C \neq 0$, we can still choose λ and β so that $c = \beta + C = 0$ and $\beta_i = -C \neq 0$ for all $i \in \mathbb{Z}$. Thus we get a simple module M with $HM = 0$ whenever ψ is nonzero. Its annihilator is $\langle H \rangle$ by Lemma 3.3. This proves (1).

When $c \neq 0$ and $o(s) = n$, we get a simple module M with $HM = M$; its annihilator must be $\langle H^n - c^n \rangle$ by Lemma 3.6. This proves (3).

When $c \neq 0$ and $o(s) = \infty$, again we get a simple module with $HM = M$. Note that even if $\psi(h) = C$, a constant (which could be zero), we can still choose λ and β so that $c = \beta + C \neq 0$ and $\beta_i = s^i c - C \neq 0$ for all $i \in \mathbb{Z}$. Thus we have a simple module M with $HM = M$ no matter what ψ is. The annihilator of this module is $\{0\}$, by Lemma 3.6. This proves (2).

Suppose now that $c \neq 0$, $o(s) = n$, and ψ is a constant C . We can still choose λ and β such that $\beta + C = c$ and $\beta_i = s^i c - C \neq 0$ for any $i \in \mathbb{Z}$, provided C is not one of the values $s^i c$, i.e., provided $C^n \neq c^n$. We get a simple module M such that $HM = M$. The annihilators must be $\langle H^n - c^n \rangle$ by Lemma 3.6. This proves (4).

It remains to prove (5) and (6). We now need to introduce modules that are not weight modules. Here is one possibility. Suppose $\psi(h) = C$ where $C \in \mathbb{C}$. Pick a positive integer n . Let M be a module with basis $\{v_i : i \in \mathbb{Z}\}$ where

$$uv_i = v_{i+1}, \quad hv_i = v_{i-n} + i\gamma v_i, \quad dv_i = (s^i - 1)Cv_{i-1}.$$

We can check that this is indeed a module:

$$\begin{aligned} huv_i &= v_{i+1-n} + (i+1)\gamma v_{i+1} = v_{i-n+1} + i\gamma v_{i+1} + \gamma v_{i+1} = u(h+\gamma)v_i; \\ dhv_i &= (s^{i-n} - 1)Cv_{i-n-1} + i\gamma(s^i - 1)Cv_{i-1} = (h+\gamma)dv_i; \\ [sud + (s-1)C]v_i &= [s^{i+1} - s]Cv_i + [s-1]Cv_i = [s^{i+1} - 1]C = duv_i. \end{aligned}$$

We now verify that this module is simple. Define the operator $T = uh - 1$. Its action on v_i is $Tv_i = i\gamma v_{i+n}$. Thus v_i is “almost” an eigenvector of T with eigenvalue $i\gamma$. In this case “almost” is good enough.

Suppose M_1 is a nonzero submodule of M . Let $x = \sum a_i v_i$ be a nonzero element in M_1 with minimal length. Then $Tx = \sum i\gamma a_i v_{i+n}$ and for any m with $a_m \neq 0$, we have $Tx - m\gamma ux = \sum (i-m)\gamma a_i v_{i+n}$, which has shorter length than x , and so must be zero. Thus $x = a_m v_m$, i.e., M_1 contains a pure basis vector v_m .

Now M can be generated by just one v_m . To get v_{m+1} we simply apply u , and to get v_{m-1} we apply h and subtract $m\gamma v_m$, getting v_{m-n} , then apply u repeatedly until we get v_{m-1} . Thus $M_1 = M$. So M is indeed a simple module.

Now if $C = 0$ and $n = 1$, then d annihilates M . By Lemma 3.3, $\text{Ann } M$ cannot be bigger than $\langle d \rangle$. Thus $\langle d \rangle$ is indeed a primitive ideal. Similarly, $\langle u \rangle$ is a primitive ideal. This proves (5).

If $C \neq 0$ and $o(s) = n$, then d^n acts as the zero scalar. Thus $\text{Ann } M$ contains $\langle d^n \rangle$. It cannot be any larger by Lemma 3.6. Thus $\langle d^n \rangle$ is a primitive ideal. Similarly, $\langle u^n \rangle$ is a primitive ideal. This proves (6). $\square$

3.5 A list of primitive ideals

Putting all these together, we get a complete list of the primitive ideals of $L(\phi, 1, s, \gamma)$ ($\gamma \neq 0$), summarized in the following table. Note that we only include primitive ideals that are not annihilators of finite-dimensional modules.

$o(s)$	ψ	primitive ideals
∞	0	$\{0\}, \langle u \rangle, \langle d \rangle$
∞	nonzero	$\{0\}, \langle H \rangle$
n	0	$\langle H \rangle, \langle u \rangle, \langle d \rangle, \langle H^n - c^n \rangle$ ($c \neq 0$)
n	$C \neq 0$	$\langle H \rangle, \langle u^n \rangle, \langle d^n \rangle, \langle H^n - c^n \rangle$ ($c \neq 0, c^n \neq C^n$)
n	nonconstant	$\langle H \rangle, \langle H^n - c^n \rangle$ ($c \neq 0$)

The preceding process of determining primitive ideals of $L(\phi, 1, s, \gamma)$ is fairly typical of our method in general. Using a central element or otherwise, we find an element that must be present in all primitive ideals. Using results similar to Lemmas 3.1 and 3.2 we hope to show that any primitive ideal larger than these minimal ideals must contain a power of u or a power of d ; usually this is enough to conclude that these large primitive ideals must be annihilators of finite-dimensional simple modules. Sometimes we will have to consider special cases that could very well involve modules that are not weight modules.

4 When L is conformal and $\gamma = 0$

In this section we assume that $\gamma = 0$ and L is conformal. Recall that this means there exists a polynomial ψ such that $s\psi(x) - \psi(rx) = \phi(x)$. We will use ψ throughout instead of ϕ .

It is very useful, especially when roots of unity are involved, to have a commutation formula between u and d^k , $k \geq 1$; see Corollary 2.4. With the base case $du - sud = s\psi(h) - \psi(rh)$, we establish by induction that

$$d^k u - s^k u d^k = [s^k \psi(h) - \psi(r^k h)] d^{k-1}.$$

There is of course a corresponding formula involving d and u^k .

As before we define H to be $ud + \psi(h)$; then $Hu = suH$ and $dH = sHd$. If M is a simple L -module, then these relations imply that either $HM = M$ or $HM = 0$; similarly, either $hM = M$ or $hM = 0$. There are thus four cases to consider: when $hM = HM = 0$, when $hM = M$ and $HM = 0$, when $hM = 0$ and $HM = M$, and finally, when $hM = HM = M$.

4.1 Homogeneous elements in ideals

Before we treat these cases individually, we state the following result. It is a somewhat more complicated analogue of Lemma 3.1.

Lemma 4.1. *Suppose $o(r) = \infty$ and x is an element in an ideal I of L . Let $x = \sum_{i \in \mathbb{Z}} x_i$ be its homogeneous decomposition (i.e., $x_i \in L_i$). Then $x_i h^{\ell(x)-1} \in I$ for all $i \in \mathbb{Z}$, where $\ell(x)$ is the length of x .*

Proof. As before, we use induction on $\ell(x)$. The claim is certainly true if $\ell(x) = 1$. Assume that $\ell(x) > 1$. Then $hx = \sum h x_i = \sum r^i x_i h$. Pick $k \in \mathbb{Z}$ such that $x_k \neq 0$. Then

$$hx - r^k x h = \sum_{i \in \mathbb{Z}} (r^i - r^k) x_i h.$$

Since $r^i \neq r^k$ unless $i = k$, we have $\ell(hx - s^k xh) = \ell(x) - 1$. By the induction hypothesis, $(r^i - r^k)x_i h h^{\ell(x)-2} \in I$; hence $x_i h^{\ell(x)-1} \in I$ for all $i \neq k$. Thus $y = \sum_{i \neq k} x_i h^{\ell(x)-1} \in I$, so $x_k h^{\ell(x)-1} = x h^{\ell(x)-1} - y \in I$ also. $\square$

Remark 4.2. The proof only uses the commutation relation between h and elements of degree i . Since H has similar commutation relations, the result is also true when h is replaced by H and r by s .

We now consider each of the four cases in turn.

4.2 When $hM = HM = 0$

We dispose of this case immediately. Both h and $ud = H - \psi(h)$ act as scalars on M , so by Lemma 2.5, M must be one-dimensional.

4.3 When $hM = M$ and $HM = 0$

Suppose first that $o(r) = n$. Then h^n commutes with u and d , and hence h^n must act as a scalar on M . This implies that h has an eigenvector; thus $hv = \lambda v$ for some nonzero $v \in M$ and $\lambda \in \mathbb{C}$. Note that $Hv = 0$, so $(ud)v = -\psi(\lambda)v$, i.e., v is also an eigenvector of ud . Thus v is a weight vector. Now recall that

$$\begin{aligned} d^k u &= s^k u d^k + [s^k \psi(h) - \psi(r^k h)] d^{k-1} \\ &= s^k [H - \psi(h)] d^{k-1} + [s^k \psi(h) - \psi(r^k h)] d^{k-1} \\ &= [s^k H - \psi(r^k h)] d^{k-1} \end{aligned}$$

for $k \geq 1$. In particular, since H acts as zero on M , we have $d^n u = -\psi(h) d^{n-1}$ as operators on M . Compare this with $ud = H - \psi(h)$, i.e., $ud^n = H d^{n-1} - \psi(h) d^{n-1}$, and we conclude that as operators on M , d^n commutes with u . Clearly d^n also commutes with h , and hence d^n acts as a scalar on M . (Similarly, u^n acts as a scalar on M .) By Lemma 2.3, M is finite-dimensional.

Thus we can concentrate on the case where r is not a root of unity. Clearly $\text{Ann } M$ contains $\langle H \rangle$, but when $\psi(h) = 0$, we can say even more. As in section 3.2, in this case $\text{Ann } M$ must actually contain u or d .

Lemma 4.3. *Suppose r is not a root of unity, $HM = 0$, $hM = M$, and $\text{Ann } M \supsetneq \langle H \rangle$. Then $\text{Ann } M$ contains a power of u or a power of d .*

When in addition $\psi(h) = 0$ and $\text{Ann } M \supsetneq \langle d \rangle$, then $\text{Ann } M$ contains u . Similarly, if $\text{Ann } M \supsetneq \langle u \rangle$, then $\text{Ann } M$ contains d .

Proof. Denote by J the ideal $\langle H \rangle$; in the special case where $\psi(h) = 0$, let J denote the ideal $\langle d \rangle$. Pick an element x in $\text{Ann } M$ that is not in J . By Lemma 4.1, $\text{Ann } M$ contains an element of the form $x_i h^k$ where $x_i \notin J$, $x_i \in L_i$. Since $hM = M$, we conclude that $x_i \in \text{Ann } M$ also. In the special case where $\psi(h) = 0$, we clearly have $i \geq 0$. In the other cases, to be definite we also assume that $i \geq 0$; the case where $i \leq 0$ is similar. Thus $\text{Ann } M$ contains an element of the form $u^i f(h)$ for some nonzero polynomial f . Among all such elements, choose one where the degree of f is as small as possible, say m . Then

$$u^i f(h)u - r^m u^{i+1} f(h) = u^{i+1} [f(rh) - r^m f(h)]$$

is also in $\text{Ann } M$. But the polynomial $f(rh) - r^m f(h)$ has smaller degree than m , hence we must have $f(rh) = r^m f(h)$. If $f(h) = \sum_{j=0}^m a_j h^j$, then this implies that $\sum_{j=0}^m (r^j - r^m) a_j h^j = 0$. Since r is not a root of unity, we have $a_j = 0$ for $j \neq m$. Thus $f(h)$ is a multiple of h^m . Since $hM = M$, we conclude that $\text{Ann } M$ contains u^i .

In the case where $\psi(h) = 0$, recall that either $uM = M$ or $uM = 0$; since $u^i \in \text{Ann } M$, we must have $uM = 0$, i.e., $u \in \text{Ann } M$.

The other cases of the lemma are treated similarly. $\square$

The lemma has immediate consequences for large primitive ideals.

Lemma 4.4. *Suppose $o(r) = \infty$, M is a simple module, $HM = 0$, and $hM = M$. If $\text{Ann } M \supsetneq \langle H \rangle$ and ψ is not zero, then M is finite-dimensional. If ψ is zero and $\text{Ann } M \supsetneq \langle u \rangle$ or $\text{Ann } M \supsetneq \langle d \rangle$, then M is one-dimensional.*

Proof. By the previous lemma, $\text{Ann } M$ contains a power of u or a power of d ; assume it is a power of d for definiteness. Then there is a nonzero element $v \in M$ such that $dv = 0$. Since $H = ud + \psi(h)$ and $HM = 0$, it is clear that $\psi(h)v = 0$ also. By Corollary 2.4, M is finite-dimensional.

If ψ is zero, then the previous lemma implies that $\text{Ann } M$ contains both u and d . Thus as operators on M , the elements u , d , and h all commute, so M must be one-dimensional. $\square$

4.4 When $hM = 0$ and $HM = M$

This case is similar to the previous one, with H and s interchanged with h and r . The conclusions, however, are different, so we will go through the details.

As before, we start by looking at the case when $o(s) = n$. The operator $\psi(h)$ acts as the scalar $\psi(0)$ on M , so in this case we have $d^k u = s^k u d^k +$

$(s^k - 1)\psi(0)d^{k-1}$. Thus $d^n u = u d^n$, i.e., d^n commutes with u (and with h) as an operator on M . Thus d^n acts as a scalar on M . Similarly, u^n acts as a scalar on M .

Note that H^n also commutes with u , d , and h , and hence acts as a scalar on M . Thus H has an eigenvector on M . Since h acts as a scalar on M (namely, as zero), we can use Lemma 2.3 to conclude that M is finite-dimensional.

We can thus concentrate on the case where s is not a root of unity. Clearly $\text{Ann } M$ contains $\langle h \rangle$. It turns out that $\text{Ann } M$ cannot be larger than $\langle h \rangle$ in this case.

Lemma 4.5. *Suppose s is not a root of unity, and M is a simple module with $HM = M$, $hM = 0$. Then $\text{Ann } M = \langle h \rangle$.*

Proof. Assume that $\text{Ann } M \supsetneq \langle h \rangle$. We will derive a contradiction. Using the same calculation as in Lemma 4.3, with H and s exchanging roles with h and r , we conclude that $\text{Ann } M$ contains a power of u or a power of d . For definiteness we assume that $\text{Ann } M$ contains d^m but not d^{m-1} , where $m \geq 1$. Then $\text{Ann } M$ also contains

$$d^m u - s^m u d^m = (s^m - 1)\psi(0)d^{m-1}.$$

If $\psi(0) \neq 0$ we reach our contradiction. If $\psi(0) = 0$, then $ud = H$ as operators on M . Thus $u^m d^m = s^{-m} H^m$, but this is also a contradiction since the $u^m d^m$ acts as zero while H^m cannot act as zero since $HM = M$. $\square$

4.5 When $hM = M$ and $HM = M$

As before, we first consider the case where both r and s are roots of unity. In this case there exists a positive integer n such that $r^n = s^n = 1$. Then h^n and H^n are both central elements, so they both act as scalars on M . Since h and H commute, they have a common eigenvector. Therefore h and $ud = H - \psi(h)$ also has a common eigenvector.

Recall that $d^k u - s^k u d^k = [s^k \psi(h) - \psi(r^k h)]d^{k-1}$, which implies that $d^n u = d^n u$. Clearly $d^n h = r^n h d^n = h d^n$, thus d^n is a central element. Hence d^n also acts as a scalar on M . Similarly, u^n also acts as a scalar on M . We can apply Lemma 2.3 and conclude that M is actually finite-dimensional.

We thus can concentrate on the case where at least one of r and s is not a root of unity. In this case we can use Lemma 4.1 and the remark following it, either through H and s , or through h and r .

Since now both h and H act nontrivially on M , it is possible that they interact in somewhat complicated ways. To help keep track of this interaction we make use of the set $\mathcal{S}(r, s) = \{(i, j) \in \mathbb{Z} \times \mathbb{Z} : r^i = s^j\}$. We will often just write $\mathcal{S}$ for $\mathcal{S}(r, s)$, since r and s are usually fixed.

Note that $\mathcal{S}$ is an additive subgroup of $\mathbb{Z} \times \mathbb{Z}$. Since we are assuming that not both of r and s are roots of unity, it is straightforward to verify that $\mathcal{S}$ is in fact generated by one element: there exists $(n, m) \in \mathcal{S}$ such that $\mathcal{S} = \{(kn, km) : k \in \mathbb{Z}\}$. We can thus divide the sets $\mathcal{S}$ into two types: those that contain pairs (i, j) where i and j have the same sign or zero, and those that contain pairs (i, j) where i and j have opposite signs. For ease of reference, we adopt the following notational convention: when $\mathcal{S}$ is of the first type, we write $\mathcal{S} = \langle (n, m) \rangle$ to indicate that $\mathcal{S}$ is generated by (n, m) where $n, m \geq 0$; when $\mathcal{S}$ is of the second type, we write $\mathcal{S} = \langle (n, -m) \rangle$ to indicate that $\mathcal{S}$ is generated by $(n, -m)$ where $n, m > 0$. Thus the type of $\mathcal{S}$ is indicated by the presence or absence of the minus sign on m .

Now let M be a simple module. Every element of $\mathcal{S}$ gives rise to an element of $\text{Ann } M$, as follows.

Suppose first that $\mathcal{S} = \langle (n, m) \rangle$ and $(i, j) \in \mathcal{S}$. The pair $(-i, -j)$ gives rise to the same element of $\text{Ann } M$, so we can assume that $i, j \geq 0$. Since $hM = M$ (and $\ker h = 0$), the operator h has an inverse h^{-1} (although the element h does not have an inverse in L). We have $h^{-1}u = r^{-1}uh^{-1}$ and $dh^{-1} = r^{-1}h^{-1}d$. Thus the operator $h^{-i}H^j$ commutes with everything in L , and hence acts as a scalar $c_{i,j}$ on M . The scalar $c_{i,j}$ must be nonzero because $hM = M$ and $HM = M$. This implies that H^j acts the same way on M as does $c_{i,j}h^i$. Therefore $\text{Ann } M$ must contain $H^j - c_{i,j}h^i$.

We define $I_{\mathcal{S}}$ as the ideal generated by all $H^j - c_{i,j}h^i$ as (i, j) ranges over the elements of $\mathcal{S}$; then $I_{\mathcal{S}} \subseteq \text{Ann } M$. Since $\mathcal{S}$ is generated by the single element (n, m) , it is straightforward to verify that if $(i, j) = (kn, km)$, then $H^j - (c_{n,m})^k h^i \in \langle H^m - c_{n,m}h^n \rangle \subseteq I_{\mathcal{S}}$. Since $H^j - c_{i,j}h^i \in I_{\mathcal{S}}$, we conclude that $c_{i,j} = (c_{n,m})^k$ and that $I_{\mathcal{S}} = \langle H^m - c_{n,m}h^n \rangle$.

Suppose now that $\mathcal{S} = \langle (n, -m) \rangle$ and $(i, -j) \in \mathcal{S}$. As before, the pair $(-i, j)$ gives rise to the same element of $\text{Ann } M$, so we can assume that $i, j > 0$. This time we conclude that $h^i H^j - c_{i,-j} \in \text{Ann } M$ for some nonzero scalar $c_{i,-j}$. A similar calculation as before shows that $c_{i,-j} = (c_{n,-m})^k$ if $(i, -j) = (kn, -km)$. If we define $I_{\mathcal{S}}$ as the ideal generated by all $h^i H^j - c_{i,-j}$ as $(i, -j)$ ranges over $\mathcal{S}$, then $I_{\mathcal{S}} = \langle h^n H^m - c_{n,-m} \rangle$.

Here are some examples. Suppose r and s are algebraically independent. Then $\mathcal{S} = \{(0, 0)\}$ and $I_{\mathcal{S}} = \{0\}$. Suppose $o(s) = m < \infty$ and $o(r) = \infty$. Then $\mathcal{S} = \{(0, km) : k \in \mathbb{Z}\}$ and $I_{\mathcal{S}} = \langle H^m - c_{0,m} \rangle$. Suppose $rs = 1$ and $o(r) = o(s) = \infty$. Then $\mathcal{S} = \{(k, -k) : k \in \mathbb{Z}\}$ and $I_{\mathcal{S}} = \langle hH - c_{1,-1} \rangle$.

Inspired by these observations, we define, for a given $\mathcal{S}$ and any $c \in \mathbb{C}^\times$, the ideal $I_c = \langle H^m - ch^n \rangle$ if $\mathcal{S} = \langle (n, m) \rangle$, and $I_c = \langle h^n H^m - c \rangle$ if $\mathcal{S} = \langle (n, -m) \rangle$. Note that if $(i, j) \in \mathcal{S}$ with $i, j \geq 0$, then $(i, j) = (kn, km)$ for some $k \geq 0$, and the element $H^j - c^k h^i$ lies in I_c . Similarly, if $(i, -j) \in \mathcal{S}$ with $i, j > 0$, then $(i, -j) = (kn, -km)$ for some $k \geq 0$, and the element $h^i H^j - c^k$ lies in I_c .

Any primitive ideal of L contains I_c for some $c \in \mathbb{C}^\times$. We need to determine which values of c actually make I_c a primitive ideal. The easy case, of course, is when $\mathcal{S} = \{(0, 0)\}$. Then $I_c = \langle 1 - c \rangle$ and so only $I_1 = \{0\}$ can be a primitive ideal.

When $\mathcal{S} \neq \{(0, 0)\}$, the situation is almost the complete opposite. It turns out that almost all nonzero values of c make I_c a primitive ideal, but as usual there are special cases where $\text{Ann } M$ has to contain more than just I_c .

Lemma 4.6. *Let M be a simple module with $hM = HM = M$. Suppose (jm, m) is the generator of $\mathcal{S}$ for some $j \geq 0$ and $m > 0$ (thus $s^m = r^{jm}$). Suppose further that $\psi(h) = Ch^j$ with $C \neq 0$. If $I_{C^m} \subseteq \text{Ann } M$, i.e., if $H^m - C^m h^{jm} \in \text{Ann } M$, then $\text{Ann } M$ also contains u^m or d^m .*

Proof. We note that $s^{-1}r^j$ is an m th root of unity. Since $du = sH - Cr^j h^j$, we get by induction that

$$d^k u^k = \prod_{i=1}^k (s^i H - r^{ij} Ch^j) = \prod_{i=1}^k s^i [H - (r^j s^{-1})^i Ch^j].$$

In particular, for $k = m$, we get that $d^m u^m$ is a nonzero multiple of $H^m - C^m h^{jm}$. Thus $d^m u^m$ acts as the zero operator on M .

Recall that

$$d^m u - s^m u d^m = [s^m \psi(h) - \psi(r^m h)] d^{m-1}.$$

When $\psi(h) = Ch^j$ and $s^m = r^{jm}$, we get $d^m u = s^m u d^m$. Thus $d^m M$ is a submodule of M , either 0 or M . Similarly, $u^m M = 0$ or $u^m M = M$. Since $d^m u^m$ acts as the zero operator on M , it cannot be the case that both $u^m M = M$ and $d^m M = M$. Therefore $\text{Ann } M$ contains either u^m or d^m . $\square$

We now look at what happens when a primitive ideal is larger than I_c , or in the case of the lemma above, larger than $\langle u^m \rangle$ or $\langle d^m \rangle$.

To do so, we need the following lemma about two-variable polynomials, analogous to Lemma 3.5. We first recall the definition of distinctive polynomials from [P1]. A set $\mathcal{T} \subseteq \mathbb{Z} \times \mathbb{Z}$ is called distinctive if $r^i s^j \neq r^{i'} s^{j'}$ for

all distinct pairs $(i, j), (i', j') \in \mathcal{T}$. A polynomial is called *distinctive* if its powers are in a distinctive set, i.e., $g(x, y) = \sum_{(i,j) \in \mathcal{T}} a_{i,j} x^i y^j$ is distinctive if $\mathcal{T}$ is a distinctive set. (We count the zero polynomial as distinctive.)

(Distinctive sets are not exotic. Note that $r^i s^j = r^{i'} s^{j'}$ iff $r^{i-i'} = s^{-(j-j')}$, i.e., iff $(i - i', -(j - j')) \in \mathcal{S}$. Thus (i, j) and (i', j') are in a distinctive set iff $(i, -j)$ and $(i', -j')$ are in different $\mathcal{S}$ -cosets. Therefore $\mathcal{T}$ is a distinctive set iff the elements $(i, -j)$, where $(i, j) \in \mathcal{T}$, are different $\mathcal{S}$ -coset representatives.)

Lemma 4.7. *Suppose $g(x, y) = \sum_{(i,j) \in \mathcal{T}} a_{i,j} x^i y^j$ is a distinctive polynomial and $g(rx, sy) = r^a s^b g(x, y)$ where $(a, b) \in \mathcal{T}$. Then $g(x, y) = cx^a y^b$ for some $c \in \mathbb{C}$.*

Proof. We have

$$\sum_{(i,j) \in \mathcal{T}} a_{i,j} (r^i s^j - r^a s^b) x^i y^j = 0.$$

Since $\mathcal{T}$ is distinctive, we have $r^i s^j \neq r^a s^b$ unless $(i, j) = (a, b)$. The conclusion follows. $\square$

Distinctive polynomials are useful because any element of L_0 is congruent modulo I_c (where $c \in \mathbb{C}^\times$) to a distinctive polynomial in h and H . To see this, suppose $g(h, H) = \sum_{(i,j) \in \mathcal{R}} a_{i,j} h^i H^j$ is an arbitrary element of L_0 , where $\mathcal{R}$ is not necessarily a distinctive set. We will show that we can reduce the set $\mathcal{R}$ until we get a distinctive set, without changing the congruence class of $g(h, H) \bmod I_c$.

So suppose $(i, j), (i', j') \in \mathcal{R}$ with $r^i s^j = r^{i'} s^{j'}$. Then $r^{i-i'} = s^{j'-j}$, so $(i - i', j' - j) \in \mathcal{S}$. Clearly it does no harm to assume that $i - i' \geq 0$. If $j' - j \geq 0$, then $H^{j'-j} - c^k h^{i-i'} \in I_c$ for some $k \geq 0$, so $H^{j'} h^{i'} \equiv c^k h^i H^j \bmod I_c$. Thus we can replace the term $h^{i'} H^{j'}$ by $c^k h^i H^j$ without changing the congruence class of $g(h, H) \bmod I_c$. On the other hand, if $j' - j < 0$, then $h^{i-i'} H^{j-j'} - c^k \in I_c$ for some $k \geq 0$. Thus $h^i H^j \equiv c^k h^{i'} H^{j'} \bmod I_c$, and we can still replace the term $h^{i'} H^{j'}$ by $c^k h^i H^j$ without changing the congruence class of $g(h, H)$. Clearly we can continue this process until we obtain a distinctive polynomial.

We use the lemma about distinctive polynomials to prove the following analogue of Lemma 3.2.

Lemma 4.8. *Let M be a simple module with $hM = HM = M$. If $\text{Ann } M \supsetneq I_c$, then $\text{Ann } M$ contains a power of u or a power of d .*

In the special case of Lemma 4.6, suppose $\text{Ann } M \supsetneq \langle u^m \rangle$. Then $\text{Ann } M$ contains d^m . Similarly, if $\text{Ann } M \supsetneq \langle d^m \rangle$, then $\text{Ann } M$ contains u^m .

Proof. By Lemma 4.1 $\text{Ann } M$ contains a homogeneous element x not in I_c . For definiteness assume that x has degree $k \geq 0$; the case $k < 0$ is similar. We can write $x = u^k f(h, H)$ where $f \in \mathbb{C}[x, y]$ is a nonzero polynomial. Since $x \notin I_c$, we can assume that f is a nonzero distinctive polynomial.

Among all elements of the form $u^k f(h, H)$ where f is nonzero and distinctive, choose one where the degree of f is minimal. Let's say the degree of f is (a, b) . Then

$$xu - r^a s^b ux = u^k f(h, H) u - r^a s^b u^{k+1} f(h, H) = u^{k+1} [f(rh, sH) - r^a s^b f(h, H)].$$

The polynomial $f(rh, sH) - r^a s^b f(h, H)$ is distinctive and has smaller degree than f , so it must be zero. By Lemma 4.7, $f(h, H)$ is a scalar multiple of $h^a H^b$, so $\text{Ann } M$ contains $u^k h^a H^b$. Since $hM = HM = M$, we conclude that $\text{Ann } M$ contains u^k .

Suppose now that we are in the special case of Lemma 4.6. Assume that $\text{Ann } M \supsetneq \langle u^m \rangle$. As above, $\text{Ann } M$ contains a homogeneous element x of degree k not in $\langle u^m \rangle$; here we can assume that $k < m$. We note that since $\text{Ann } M$ contains u^m , it also contains $d^{m-k} u^m = d^{m-k} u^{m-k} u^k$. Here $d^{m-k} u^{m-k} = \prod_{i=1}^{m-k} s^i [H - (r^j s^{-1})^i C h^j]$ is a polynomial whose H -degree is $m - k$. Thus we can write $x = u^k f(h, H)$ where f is a polynomial whose H -degree is smaller than $m - k$. Therefore $\text{Ann } M$ contains the element $d^k x = d^k u^k f(h, H) = g(h, H)$, a polynomial whose H -degree is at most $m - 1$. Since $\mathcal{S} = \{i(jm, m) : i \in \mathbb{Z}\}$, the polynomial g is distinctive. Thus $\text{Ann } M$ contains an element of the form $d^i g(h, H)$ where g is nonzero and distinctive. By the calculation above, $\text{Ann } M$ contains a power of d . Since $d^m M = 0$ or $d^m M = M$, we conclude that $d^m \in \text{Ann } M$, as required. $\square$

In most cases, if $\text{Ann } M$ contains a power of u or a power of d , then M must be finite-dimensional.

Lemma 4.9. *Let M be a simple module with $hM = HM = M$. Suppose $u^m \in \text{Ann } M$ but $u^{m-1} \notin \text{Ann } M$, $m \geq 1$. Then ψ is not the zero polynomial and M is finite-dimensional, except in the special case of Lemma 4.6. The same conclusion holds when $d^m \in \text{Ann } M$ and $d^{m-1} \notin \text{Ann } M$.*

In the special case of Lemma 4.6, if $\text{Ann } M$ contains both u^m and d^m , then M is finite-dimensional.

Proof. If ψ is the zero polynomial, then $du = sud$. Thus uM is a submodule and hence must be either 0 or M . (Similarly, either $dM = 0$ or $dM = M$.) Since $u^m M = 0$, we must have $uM = 0$. Thus $ud = H$ acts as the zero operator on M , contradicting $HM = M$.

So assume now that ψ is nonzero. Recall that

$$d^m u - s^m u d^m = [s^m \psi(h) - \psi(r^m h)] d^{m-1}.$$

If $\psi(h) = \sum_i a_i h^i$, then $s^m \psi(h) - \psi(r^m h) = \sum_i a_i (s^m - r^{im}) h^i$, which is nonzero unless $s^m = r^{jm}$ for some $j \geq 0$ and $a_i = 0$ for $i \neq j$, i.e., unless we are in the special situation of Lemma 4.6. (Note that the equality $s^m = r^{jm}$ means that r cannot be a root of unity, since that would force s to be a root of unity also, and we have excluded this possibility. Thus $s^m = r^{jm}$ is true only for at most one value of j .) We obtain the conclusion from Corollary 2.4.

Suppose now we are in the situation of Lemma 4.6. The proof in this case is quite similar to the proof of the last case of Lemma 3.7. Choose an element $v \in M$ such that $dv = 0$ (such an element exists since $d^m \in \text{Ann } M$). Define $v_i = u^i v$ for $i \geq 0$; since $u^m \in \text{Ann } M$, we have $u_{m+i} = 0$ for $i \geq 0$.

Now $0 = udv_0 = (H - Ch^j)v_0$, so $Hv_0 = Ch^j v_0$. Recall that as an operator on M , the operator h has an inverse h^{-1} (although the element h is not invertible in L); thus $Hh^{-j}v_0 = Cv_0$. Since $Hh^{-j}u = sr^{-j}uHh^{-j}$, we see that $Hh^{-j}v_i = (sr^{-j})^i v_i$ for $0 \leq i \leq m-1$. Each v_i is an eigenvector of the operator Hh^{-j} , with distinct eigenvalues.

Since M is simple, there exists $x \in L$ such that $xhv_0 = v_0$. Since $dhv_0 = 0$, we can assume that x can be written as $\sum_i p_i(h)u^i$, so

$$\sum_i p_i(h)u^i hv_0 = \sum_i r^{-i} p_i(h) h u^i v_0 = \sum_i r^{-i} p_i(h) h v_i = v_0.$$

For each i , $r^{-i} p_i(h) h v_i$ is an eigenvector of Hh^{-j} with eigenvalue $(sr^{-j})^i$. They form a linearly independent set. Thus we have $p_i(h) = 0$ for $i > 0$. Therefore $(p_0(h)h - 1)v_0 = 0$, i.e., there exists a nontrivial polynomial in h that annihilates v_0 . Thus we can apply Lemma 2.3 to conclude that M is finite-dimensional. $\square$

4.6 Proving primitivity

It remains to show that the ideals $I_c, \langle h \rangle, \langle H \rangle, \langle u \rangle, \langle d \rangle, \langle u^m \rangle, \langle d^m \rangle$ are all primitive ideals under the right conditions.

Lemma 4.10. *Suppose L is conformal, with $\gamma = 0$. Then the following ideals are primitive:*

1. $\langle h \rangle$ if $o(s) = \infty$;
2. $\langle H \rangle$ if $o(r) = \infty$ and ψ is not identically zero;

3. $\langle u \rangle$ and $\langle d \rangle$ if $o(r) = \infty$ and ψ is identically zero;
4. $\{0\}$ if $\mathcal{S}(r, s) = \{(0, 0)\}$;
5. $\langle h^n H^m - c \rangle$ if $c \in \mathbb{C}^\times$ and $\mathcal{S}(r, s) = \langle (n, -m) \rangle$ (where $n, m > 0$);
6. $\langle H^m - ch^n \rangle$, where $c \in \mathbb{C}^\times$, $\mathcal{S}(r, s) = \langle (n, m) \rangle$ (where one of n or m is nonzero), and we are not in the case of Lemma 4.6;
7. $\langle u^m \rangle$ and $\langle d^m \rangle$ if we are in the case of Lemma 4.6, i.e., if $\psi(h) = Ch^j$ for some $C \neq 0$ and $j \geq 0$, and $\mathcal{S}(r, s) = \langle (jm, m) \rangle$ for some $m > 0$.

Proof. We again make use of the universal weight module $W(\lambda, \beta)$ described in section 2.1. In this case, it is more convenient to use the parameters λ and $\mu = \beta + \psi(\lambda)$ instead of λ and β . We have $hv_i = r^i \lambda v_i$ and $Hv_i = s^i \mu v_i$, hence $udv_i = (H - \psi(h))v_i = [s^i \mu - \psi(r^i \lambda)]v_i$. The module $W(\lambda, \mu)$ is simple if the weights $(r^i \lambda, s^i \mu)$ are distinct and $s^i \mu - \psi(r^i \lambda) \neq 0$ for all $i \in \mathbb{Z}$.

Suppose now that $o(s) = \infty$. Set $\lambda = 0$. Then the weights $(r^i \lambda, s^i \mu) = (0, s^i \mu)$ are certainly distinct if $\mu \neq 0$, and we can choose a nonzero value of μ such that $s^i \mu - \psi(0) \neq 0$ for all $i \in \mathbb{Z}$. Thus $W(0, \mu)$ is simple; its annihilator is $\langle h \rangle$ by Lemma 4.5. This proves (1).

Suppose next that $o(r) = \infty$ and ψ is not identically zero. Set $\mu = 0$. The weights $(r^i \lambda, s^i \mu) = (r^i \lambda, 0)$ are distinct if $\lambda \neq 0$. In this case, $s^i \mu - \psi(r^i \lambda)$ is equal to $-\psi(r^i \lambda)$. Since ψ is not identically zero, there are only finitely many values x such that $\psi(x) = 0$; thus there are only countably many values of λ such that $\psi(r^i \lambda) = 0$ for some $i \in \mathbb{Z}$. If we choose a nonzero value of λ outside these countably many special values, then $\psi(r^i \lambda) \neq 0$ for all $i \in \mathbb{Z}$. For this value of λ , $W(\lambda, 0)$ is simple. Its annihilator must be $\langle H \rangle$ by Lemma 4.4. This proves (2).

Suppose now that $\mathcal{S}(r, s) = \{(n, 0)\}$. Then s is not a root of unity, so the weights $(r^i \lambda, s^i \mu)$ are distinct when $\mu \neq 0$. Now if $n \neq 0$, then for any $c \in \mathbb{C}^\times$, choose λ so that $\lambda^n = 1/c$. If $n = 0$, we can choose any nonzero value for λ ; in this case c has to be 1. Then h^n acts as the scalar $\lambda^n = c$. Furthermore, there are only countably many values of $s^{-i} \psi(r^i \lambda)$ as i ranges over $\mathbb{Z}$, so if we choose μ to be different from these countably many values, then we have $s^i \mu - \psi(r^i \lambda) \neq 0$ for all $i \in \mathbb{Z}$. Then $W(\lambda, \mu)$ is simple, and its annihilator must be I_c (or I_1 when $n = m = 0$) by Lemmas 4.8 and 4.9. This proves (1) and parts of (6).

Suppose next that $\mathcal{S}(r, s) = \langle (n, -m) \rangle$. Then $h^n H^m v_i = r^n s^m \lambda^n \mu^m v_i = \lambda^n \mu^m v_i$, i.e., $h^n H^m$ acts as the scalar $\lambda^n \mu^m$ on $W(\lambda, \mu)$. I claim that for any $c \in \mathbb{C}^\times$, we can find values for λ and μ such that $\lambda^n \mu^m = c$, $(r^i \lambda, s^i \mu)$ are all distinct, and $s^i \mu - \psi(r^i \lambda) \neq 0$ for all $i \in \mathbb{Z}$. This would imply that

$W(\lambda, \mu)$ is simple. Its annihilator contains I_c , and by Lemma 4.9 it cannot be larger than I_c .

To prove the claim, set $\mu^m = c/\lambda^n$. Then the equation $s^i\mu - \psi(r^i\lambda) = 0$ implies that $s^{im}c = \lambda^n\psi(r^i\lambda)^m$. If ψ is the zero polynomial, then there is no solution for any $i \in \mathbb{Z}$. Otherwise, for each $i \in \mathbb{Z}$, there are only finitely many values of λ satisfying the equation (recall that $n > 0$, so $\lambda^n\psi(r^i\lambda)^m$ is a nontrivial polynomial in λ). Thus there are at most countably many values of λ such that $s^{im}c = \lambda^n\psi(r^i\lambda)^m$ for some $i \in \mathbb{Z}$. If we choose a nonzero value of λ different from any of these countably many values, we get $s^{im}c/\lambda^n \neq \psi(r^i\lambda)^m$ for all $i \in \mathbb{Z}$, and hence $s^i\mu - \psi(r^i\lambda) \neq 0$ for all $i \in \mathbb{Z}$. This proves the claim and finishes the proof of (5).

Suppose next that $\mathcal{S}(r, s) = \langle (n, m) \rangle$ with $m \neq 0$. Then $H^m v_i = s^m \mu^m v_i = r^n \mu^m v_i = (\mu^m/\lambda^n) r^n \lambda^n v_i = (\mu^m/\lambda^n) h^n v_i$, i.e., $H^m - (\mu^m/\lambda^n) h^n$ is in $\text{Ann } W(\lambda, \mu)$. I now claim that, except in the case of Lemma 4.6, for any $c \in \mathbb{C}^\times$ we can always find values of λ and μ such that $c = \mu^m/\lambda^n$, $(r^i\lambda, s^i\mu)$ are all distinct, and $s^i\mu - \psi(r^i\lambda) \neq 0$ for all $i \in \mathbb{Z}$. As before, this would imply that $W(\lambda, \mu)$ is simple; its annihilator contains I_c , and by Lemma 4.9 it cannot be larger than I_c .

To prove the claim, we set $\mu^m = c\lambda^n$. Then the equation $s^i\mu - \psi(r^i\lambda) = 0$ implies that $s^{im}c\lambda^n = \psi(r^i\lambda)^m$. This is a nontrivial polynomial equation in λ , unless $\psi(h) = Ch^j$ for some $j \geq 0$, $n = jm$, and $r^{ijm}C^m = s^{im}c$ (and note that the last equality implies $C^m = c$ since $r^{jm} = s^m$). Thus outside the special case of Lemma 4.6, there are at most countably many values of λ such that $s^i\mu - \psi(r^i\lambda) = 0$ for some $i \in \mathbb{Z}$. All we have to do now is pick a nonzero value of λ outside this countably many set of values; as before, this suffices to prove the claim. We have now proven (6).

We now tackle the special case of Lemma 4.6. Recall that in this case we have $r^n = s^m$, $n = jm$, $\psi(h) = Ch^j$, and $o(r^j s^{-1}) = m$. Write θ for $r^j s^{-1}$. We construct an infinite-dimensional simple module whose annihilator is $\langle d^m \rangle$; we can show that $\langle u^m \rangle$ is primitive in a similar fashion.

The construction proceeds as follows. (Note that it works even when $j = 0$.) The module M has basis $\{v_i : i \in \mathbb{Z}\}$, and the action of L on M is given by

$$uv_i = v_{i+1}, \quad dv_i = Cs^i(1 - \theta^i)v_{i-n-1}, \quad hv_i = r^{i+(n-m)/2}v_{i-m},$$

note that we might need a square root of r . It is not hard to check that $huv_i = ruhv_i$ for all $i \in \mathbb{Z}$. Verifying the other relations is not as straightforward. We have

$$dhv_i = r^{i+(n-m)/2}dv_{i-m} = Cr^{i+(n-m)/2}s^{i-m}(1 - \theta^{i-m})v_{i-m-n-1},$$

while

$$rhdv_i = rCs^i(1 - \theta^i)hv_{i-n-1} = Cr^{i-n+(n-m)/2}s^i(1 - \theta^i)v_{i-n-1-m}.$$

Since $\theta^m = 1$ and $r^n = s^m$, the two expressions are equal, proving that $dhv_i = rhdv_i$ for all $i \in \mathbb{Z}$.

To verify the last relation, we first calculate that $h^jv_i = r^{ij}v_{i-n}$. In the algebra L , we have $du - sud = (s - r^j)Ch^j$. Now

$$(du - sud)v_i = Cs^{i+1}[(1 - \theta^{i+1}) - (1 - \theta^i)]v_{i-n} = Cs^{i+1}\theta^i(1 - \theta)v_{i-n},$$

while

$$(s - r^j)Ch^jv_i = (s - r^j)Cr^{ij}v_{i-n} = C(s - s\theta)(s\theta)^iv_{i-n} = Cs^{i+1}\theta^i(1 - \theta)v_{i-n},$$

thus verifying that M is indeed an L -module.

It is easy to show that M is simple. Each v_i is an eigenvector of u^mh with eigenvalue $r^{(n-m)/2}r^i$. These eigenvalues are distinct (recall that r is not a root of unity), implying that M is simple. Note that $d^m \in \text{Ann } M$; by Lemma 4.9 we must have $\text{Ann } M = \langle d^m \rangle$. This proves (7).

We now note that the construction above works when $C = 0$ (hence $\psi = 0$) and $n = m = 1$. The action of L becomes

$$uv_i = v_{i+1}, \quad dv_i = 0, \quad hv_i = r^i v_{i-1}.$$

It is straightforward to check that with this action of L , M is still an L -module no matter the values of r and s , as long as ψ is identically zero. When r is not a root of unity, the proof above shows that M is simple. Also as above, its annihilator is $\langle d \rangle$. Thus $\langle d \rangle$ is primitive. Similarly, $\langle u \rangle$ is primitive. This proves (3). $\square$

4.7 A list of primitive ideals

We now summarize our results and provide a list of primitive ideals of L when L is conformal. As before we include only primitive ideals that are annihilators of infinite-dimensional modules. In the first table below, we list the primitive ideals when r or s is a root of unity.

$o(r)$	$o(s)$	ψ	primitive ideals
n	m	any	—
n	∞	any	$\langle h \rangle, \langle h^n - c \rangle$ ($c \in \mathbb{C}^\times$)
∞	m	0	$\langle u \rangle, \langle d \rangle, \langle H^m - c \rangle$ ($c \in \mathbb{C}^\times$)
∞	m	$C \neq 0$	$\langle u^m \rangle, \langle d^m \rangle, \langle H \rangle, \langle H^m - c^m \rangle$ ($c \in \mathbb{C}^\times, c^m \neq C^m$)
∞	m	nonconstant	$\langle H \rangle, \langle H^m - c \rangle$ ($c \in \mathbb{C}^\times$)

In the second table, we list primitive ideals when both r and s are not roots of unity. The results are expressed in terms of the generator of $\mathcal{S}(r, s)$. In the table, n , m , and j are all positive integers.

generator	ψ	primitive ideals
$(0, 0)$	0	$\{0\}, \langle h \rangle, \langle u \rangle, \langle d \rangle$
$(0, 0)$	nonzero	$\{0\}, \langle h \rangle, \langle H \rangle$
$(n, -m)$	0	$\langle h \rangle, \langle u \rangle, \langle d \rangle, \langle h^n H^m - c \rangle \ (c \in \mathbb{C}^\times)$
$(n, -m)$	nonzero	$\langle h \rangle, \langle H \rangle, \langle h^n H^m - c \rangle \ (c \in \mathbb{C}^\times)$
(jm, m)	$Ch^j \ (C \neq 0)$	$\langle h \rangle, \langle H \rangle, \langle u^m \rangle, \langle d^m \rangle, \langle H^m - ch^{jm} \rangle \ (c \in \mathbb{C}^\times, c \neq C^m)$
(jm, m)	0	$\langle h \rangle, \langle u \rangle, \langle d \rangle, \langle H^m - ch^{jm} \rangle \ (c \in \mathbb{C}^\times)$
(jm, m)	$\neq 0, \neq Ch^j$	$\langle h \rangle, \langle H \rangle, \langle H^m - ch^{jm} \rangle \ (c \in \mathbb{C}^\times)$
$(n, m), n \neq jm$	0	$\langle h \rangle, \langle u \rangle, \langle d \rangle, \langle H^m - ch^n \rangle \ (c \in \mathbb{C}^\times)$
$(n, m), n \neq jm$	nonzero	$\langle h \rangle, \langle H \rangle, \langle H^m - ch^n \rangle \ (c \in \mathbb{C}^\times)$

5 When L is not conformal, $\gamma = 0$

In this section we assume that L is not conformal and $\gamma = 0$. If $\phi(h) = \sum_i a_i h^i$, recall our assumption means $s = r^j$ and $a_j \neq 0$ for some $j \geq 0$. If r is not a root of unity, then there is only one such value of j , but if r is a root of unity, then there could be more than one value of j .

We now establish some notation. Suppose first that r is a root of unity, with $o(r) = n$. We fix j so that $s = r^j$ and $0 \leq j \leq n - 1$. The polynomial ϕ can then be written as a sum of a conformal part ϕ_0 and a nonconformal part ϕ_1 , where

$$\phi_0(h) = \sum_{i \not\equiv j} a_i h^i, \quad \phi_1(h) = \sum_{i \equiv j} a_i h^i,$$

(the congruences are mod n). Note that ϕ_0 is conformal, while ϕ_1 can be written as $\phi_1(h) = sh^j \tilde{\phi}(h^n)$ for some polynomial $\tilde{\phi}$. (We use the constant factor s in order to make subsequent formulas easier to work with.) If ψ is a conformal polynomial corresponding to ϕ_0 , then $\phi(h) = s\psi(h) - \psi(rh) + sh^j \tilde{\phi}(h^n)$.

The case $n = 1$ (i.e., $r = s = 1$) perhaps deserves a special mention. Here $\phi_0(h)$ is the zero polynomial, and $\tilde{\phi}(h^n) = \tilde{\phi}(h) = s^{-1}\phi(h)$. Thus we can take $\psi(h)$ to be zero, and all the action takes place in the nonconformal part.

The situation is a bit simpler when r is not a root of unity. We still split ϕ into a conformal part and a nonconformal part, but the nonconformal part is just a monomial: $\phi_1(h) = sCh^j$ for some $C \neq 0$.

Note that this second case can be subsumed into the previous case when $\tilde{\phi}(h) = C$. Thus we can include the second case in the first, even though h^n doesn't make sense when r and s are not roots of unity.

Indeed, the two cases can be made even more similar. Suppose again that $o(r) = n$. Since $hu = ruh$, we have $h^nu = r^nuh^n = uh^n$. Similarly, $dh^n = h^nd$. Thus h^n is a central element, so $\tilde{\phi}(h^n)$ also commutes with everything in L . It follows that if M is a simple module, then $\tilde{\phi}(h^n)$ acts as a scalar $C = C(M)$. If $C = 0$, then the nonconformal part $\phi_1(h)$ acts as the zero operator on M , and thus we are in the same situation as the conformal case. Looking at the first line of the first table in section 4.7, we conclude that in this case M must be finite-dimensional. Therefore we can assume from now on that the scalar C is nonzero; we can then write $\phi(h) = s\psi(h) - \psi(rh) + sCh^j$ as operators on M . This is just like the case where $o(r) = \infty$.

In either case, just as in the conformal case, we define H to be $ud + \psi(h)$. A straightforward calculation then shows that $Hu = su(H + \tilde{\phi}(h^n)h^j)$ and $dH = s(H + \tilde{\phi}(h^n)h^j)d$. Then $Hx_i = s^i x_i(H + i\tilde{\phi}(h^n)h^j)$ for $x_i \in L_i$. We can replace $\tilde{\phi}(h^n)$ by the nonzero constant C if the terms are viewed as operators on a simple module M .

5.1 Homogeneous elements in ideals

We have the following lemma about homogeneous elements in ideals, the analogue of Lemmas 3.1 and 4.1. It is unavoidably more complicated in the case $o(r) = n$.

Lemma 5.1. *Let I be an ideal of L . If $o(r) = n$, assume also that $\tilde{\phi}(h^n) - C \in I$, where $C \neq 0$. Suppose $x \in I$ and $x = \sum_{i \in \mathbb{Z}} x_i$ (where $x_i \in L_i$) is the homogeneous decomposition of x . Then there exists an integer $m \geq 0$ such that $x_i h^m \in I$ for all $i \in \mathbb{Z}$.*

Proof. If $o(r) = \infty$, this is simply Lemma 4.1. Suppose now that $o(r) = n$. We again use induction on the length of x . For a given $k \in \mathbb{Z}$ with $x_k \neq 0$, we have

$$hx - r^k xh = \sum_{i \in \mathbb{Z}} (r^i - r^k) x_i h.$$

The element $hx - r^k xh$ thus has smaller length than x . Then for all $i \in \mathbb{Z}$ such that $r^i \neq r^k$, we have $x_i h^m \in I$ for some $m \geq 0$. Therefore the element

$y = \sum_{\{i:r^i=r^k\}} x_i h^m$ is also in I .

It follows that $Hy = \sum_{\{i:r^i=r^k\}} s^i x_i (H + i\tilde{\phi}(h^n)h^j)h^m$ is also in I . Since $\tilde{\phi}(h^n) - C$ is an element of I , we conclude that $z = \sum_{\{i:r^i=r^k\}} s^i x_i (H + iCh^j)h^m \in I$. Now recall that $r^i = r^k$ implies that $s^i = s^k$. Thus

$$\begin{aligned} & z - s^k y H - k s^k C y h^j \\ &= \sum_{\{i:r^i=r^k\}} [s^i x_i H h^m + i s^i C x_i h^{j+m}] - \sum_{\{i:r^i=r^k\}} [s^k x_i h^m H + k s^k C x_i h^{m+j}] \\ &= \sum_{\{i:r^i=r^k\}} s^k (i - k) C x_i h^{m+j}. \end{aligned}$$

This element has smaller length than y , so by the induction hypothesis, for each $i \neq k$, the homogeneous element $x_i h^{m'}$ is in I . It follows that $x_k h^{m'}$ is also in I . $\square$

Now let M be a simple module. As before, either $hM = 0$ or $hM = M$. We first look at the situation where $hM = 0$.

5.2 The case $hM = 0$

In this case h acts as the zero operator on M . Recall that the nonconformal part of $\phi(h)$ can be written as $\phi_1(h) = sCh^j$, where $C \neq 0$. Thus if $j > 0$, then $\phi_1(h)$ also acts as the zero operator on M . In this case we have exactly the same situation as the conformal case: if s is not a root of unity, then $\text{Ann } M$ cannot be larger than $\langle h \rangle$, but if s is a root of unity, then M must be finite-dimensional.

Suppose now that $j = 0$. Then $s = 1$ and $\phi(0) = C$; also, $Hu = u(H+C)$ and $dH = (H+C)d$. These relations are exactly analogous to the relations between h and u (and between h and d) in the case $r = 1, \gamma \neq 0$, with H taking the role of h and C taking the role of γ . In particular, Lemmas 3.1 and 3.2 are valid; that is, if $\text{Ann } M \supsetneq \langle h \rangle$, then $\text{Ann } M$ contains a power of u or a power of d . Let's say $\text{Ann } M$ contains d^k but not d^{k-1} for some $k > 0$. Since $du = ud + C$ as operators on M , we have

$$d^k u = u d^k + k C d^{k-1};$$

this is a contradiction since it implies that $C d^{k-1}$ annihilates M . Thus in this case, $\text{Ann } M$ cannot be larger than just $\langle h \rangle$.

5.3 The case $hM = M$

If $o(r) = n$, recall that h^n acts as a scalar on M . Thus any primitive ideal must contain $\langle h^n - c \rangle$ for some $c \in \mathbb{C}$. We are assuming that $\tilde{\phi}(c) = C \neq 0$; since $hM = M$, it is also true that $c \neq 0$. (On the other hand, if $o(r) = \infty$, we have no obvious information on what a primitive ideal must contain.)

We need the following result on polynomials of two variables. It is a version of Lemmas 3.5 and 4.7, but unfortunately it is rather more complicated.

Lemma 5.2. *Suppose $r^j = s$ and $f(x, y) = \sum_{i=0}^b g_i(x)y^i$ is a two-variable polynomial of degree (a, b) , where $\deg g_i(x) < o(r)$ for $0 \leq i \leq b$. If*

$$f(rx, sCx^j + sy) = r^a s^b f(x, y)$$

for some $C \neq 0$, then $f(x, y)$ is a constant multiple of x^a .

Proof. We first want to show that $b = 0$. Assume for a contradiction that $b > 0$. Then we can write

$$f(x, y) = g_b(x)y^b + g_{b-1}(x)y^{b-1} + \dots$$

where $g_b(x)$ has degree a . Then

$$\begin{aligned} f(rx, sCx^j + sy) &= g_b(rx)(sCx^j + sy)^b + g_{b-1}(rx)(sCx^j + sy)^{b-1} + \dots \\ &= s^b g_b(rx)y^b + s^b g_b(rx)bCx^j y^{b-1} + s^{b-1} g_{b-1}(rx)y^{b-1} + \dots \\ &= s^b g_b(rx)y^b + [bCs^b x^j g_b(rx) + s^{b-1} g_{b-1}(rx)]y^{b-1} + \dots \end{aligned}$$

This is equal to

$$r^a s^b g_b(x)y^b + r^a s^b g_{b-1}(x)y^{b-1} + \dots$$

Equating the coefficients of y^b and of y^{b-1} , we get

$$\begin{aligned} g_b(rx) &= r^a g_b(x), \\ bCs^b x^j g_b(rx) + s^{b-1} g_{b-1}(rx) &= r^a s^b g_{b-1}(x). \end{aligned}$$

Simplifying the second equation gives us

$$bCr^{a+j}x^j g_b(x) = r^{a+j}g_{b-1}(x) - g_{b-1}(rx).$$

The left-hand side is a polynomial of degree $a+j$ since g_b has degree a . Now look at the right-hand side. If $g_{b-1}(x) = \sum c_i x^i$, then

$$r^{a+j}g_{b-1}(x) - g_{b-1}(rx) = \sum c_i (r^{a+j} - r^i)x^i,$$

so the coefficient of x^{a+j} with is 0. We have the required contradiction, thus indeed $b = 0$.

We therefore conclude that $f(x, y) = g(x)$. Suppose $g(x) = \sum_{i=0}^a c_i x^i$; recall that $a < o(r)$. We require that $g(rx) = r^a g(x)$, but this implies that $\sum c_i (r^i - r^a) x^i = 0$, thus showing that $c_i = 0$ for $i \neq a$. We conclude that $g(x) = c_a x^a$, as required. $\square$

As usual, we use this lemma to prove that large primitive ideals contain a power of u or a power of d .

Lemma 5.3. *Suppose M is simple. If $o(r) = n$ and $\text{Ann } M \supsetneq \langle h^n - c \rangle$ (where $c \neq 0$), then $\text{Ann } M$ contains a power of u or a power of d . Similarly, if $o(r) = \infty$ and $\text{Ann } M \neq \{0\}$, then $\text{Ann } M$ contains a power of u or a power of d .*

Proof. Write I for the zero ideal if $o(r) = \infty$ and for $\langle h^n - c \rangle$ if $o(r) = n$. Suppose $\text{Ann } M \supsetneq I$. By Lemma 5.1, $\text{Ann } M$ contains a homogeneous element not in I . It does no harm to assume that this homogeneous element has nonnegative degree. Thus $\text{Ann } M$ contains elements of the form $x = u^k f(h, H)$, where f is a nonzero polynomial that can be written as $f(h, H) = \sum_i g_i(h) H^i$ with the degree of $g_i(x)$ being less than $o(r)$. Choose $x = u^k f(h, H)$ to be such an element where the degree (a, b) of the polynomial f is as small as possible. Then

$$xu - r^a s^b ux = u^{k+1} [f(rh, sH + sCh^j) - r^a s^b f(h, H)]$$

is also in $\text{Ann } M$, and the degree of the the polynomial $f(rh, sH + sCh^j) - r^a s^b f(h, H)$ is smaller than (a, b) . Thus it must be zero. By Lemma 5.2, $f(h, H)$ must be a (nonzero) multiple of h^a . Since $hM = M$, we conclude that $u^k \in \text{Ann } M$, as required. $\square$

Lemma 5.4. *Let M be a simple module and suppose $\text{Ann } M$ contains a power of u or a power of d . Then M is finite-dimensional.*

Proof. Let's say $\text{Ann } M$ contains d^k but not d^{k-1} . Since $du = sud + \phi_0(h) + sCh^j$, we have

$$d^k u - s^k u d^k = \left[\sum_{i=0}^{k-1} s^i \phi_0(r^{k-i-1} h) + k s^k Ch^j \right] d^{k-1}.$$

The polynomial in square brackets is nonzero since none of the monomials in ϕ_0 can cancel out h^j . By Corollary 2.4, M is finite-dimensional. $\square$

It remains to show that $\{0\}$ is a primitive ideal if $o(r) = \infty$; also that $\langle h^n - c \rangle$ is a primitive ideal when $o(r) = n$.

Lemma 5.5. *Suppose L is not conformal, $s = r^j$, and r is not a root of unity. Then $\{0\}$ is a primitive ideal.*

On the other hand, suppose $o(r) = n$ and $c \in \mathbb{C}^\times$ satisfies $\tilde{\phi}(c) \neq 0$. Then the ideal $\langle h^n - c \rangle$ is a primitive ideal.

Suppose further that $j = 0$ and $\tilde{\phi}(0) \neq 0$. Then $\langle h \rangle$ is a primitive ideal.

Proof. As usual, we use the universal weight modules $W(\lambda, \beta)$. As before, it is more convenient to use the parametrization (λ, μ) where $\mu = \beta + \psi(\lambda)$ is the H -eigenvalue of the vector v_0 . Since $Hu = su(H + Ch^j)$ (where $C = \tilde{\phi}(c)$ if $o(r) = n$; if $o(r) = \infty$, C is simply the coefficient of h^j in $\phi_1(h)$), we can figure out that $Hv_i = s^i(\mu + iC\lambda^i)v_i$ for all $i \in \mathbb{Z}$. Thus

$$udv_i = [s^i(\mu + iC\lambda^i) - \psi(r^i\lambda)]v_i.$$

Note that each v_i is an eigenvector of H with distinct eigenvalues; thus $W(\lambda, \mu)$ is simple if $\beta_i = s^i(\mu + iC\lambda^i) - \psi(r^i\lambda)$ is nonzero for all $i \in \mathbb{Z}$.

Now for any given value of λ , there are only countably many values of $s^{-i}\psi(r^i\lambda) - iC\lambda^i$ as i ranges over $\mathbb{Z}$. If we pick μ outside these values, then $\beta_i \neq 0$ for all $i \in \mathbb{Z}$ and $W(\lambda, \mu)$ would be simple. We now see what happens for various values of λ .

If $\lambda = 0$, then h acts as the zero operator on $W(\lambda, \mu)$. If in addition $j = 0$ and $\tilde{\phi}(0) \neq 0$, then the annihilator of $W(\lambda, \mu)$ cannot be larger than $\langle h \rangle$, as discussed in section 5.2. Thus $\langle h \rangle$ is indeed a primitive ideal in this situation.

If $\lambda \neq 0$, then h does not act as the zero operator, hence $hW(\lambda, \mu) = W(\lambda, \mu)$. If in addition r is not a root of unity, then the annihilator of $W(\lambda, \mu)$ cannot be larger than $\{0\}$, by Lemmas 5.3 and 5.4, so $\{0\}$ is indeed a primitive ideal.

If $o(r) = n$, then for any value $c \in \mathbb{C}^\times$ such that $\tilde{\phi}(c) \neq 0$, we can pick $\lambda \in \mathbb{C}$ with $\lambda^n = c$ (note that $\lambda \neq 0$); after that, we pick μ as above to make $W(\lambda, \mu)$ a simple module. Then h^n acts as the scalar c on $W(\lambda, \mu)$. The annihilator of $W(\lambda, \mu)$ is just $\langle h^n - c \rangle$ by Lemmas 5.3 and 5.4. $\square$

5.4 A list of primitive ideals

We can put all these results together to generate an explicit list of the primitive ideals of L when L is not conformal. Recall that $s = r^j$. If $o(r) = n$, we write $\phi(h) = \phi_0(h) + s\tilde{\phi}(h^n)h^j$ where $\phi_0(h) = \sum_{i \neq j} a_i h^i$. As

before, we list primitive ideals that are annihilators of infinite-dimensional modules.

j	$o(r)$	$p(0)$	primitive ideals
any	∞	n/a	$\{0\}, \langle h \rangle$
0	n	0	$\langle h^n - c \rangle, c \in \mathbb{C}^\times, \tilde{\phi}(c) \neq 0$
0	n	nonzero	$\langle h \rangle, \langle h^n - c \rangle, c \in \mathbb{C}^\times, \tilde{\phi}(c) \neq 0$
> 0	n	any	$\langle h^n - c \rangle, c \in \mathbb{C}^\times, \tilde{\phi}(c) \neq 0$

References

- [BR] G. Benkart and T. Roby, Down-up algebras, *J. Algebra* **209** (1998), 305–344; Addendum, **213** (1999), 378.
- [CL] P. Carvalho and S. Lopes, Automorphisms of generalized down-up algebras, *Comm. Algebra* **37** (2009), 1622–1646.
- [CS] T. Cassidy and B. Shelton, Basic properties of generalized down-up algebras, *J. Algebra* **279** (2004), 402–421.
- [D] J. Dixmier, *Enveloping Algebras*, American Mathematical Society, Providence, RI (1996).
- [J] D. Jordan, Down-up algebras and ambiskew polynomial rings, *J. Algebra* **228** (2000), 311–346.
- [LB] L. Le Bruyn, Conformal $\mathfrak{sl}_2$ enveloping algebras, *Comm. Algebra* **23** (1995), 1325–1362.
- [MR] J. McConnell and J. Robson, *Noncommutative Noetherian Rings*, John Wiley and Sons, New York, 1987.
- [P1] I. Praton, Primitive ideals of noetherian down-up algebras, *Comm. Algebra* **32** (2004), 443–471.
- [P2] I. Praton, Simple weight modules of non-noetherian generalized down-up algebras, *Comm. Algebra*, **35** (2007), 325–337.
- [P3] I. Praton, Simple Modules and Primitive Ideals of Non-Noetherian Generalized Down-Up Algebras, *Comm. Algebra* **37** (2009), 811–839.

- [PM] I. Praton and S. May, Primitive ideals of non-noetherian down-up algebras, *Comm. Algebra* **33** (2005), 605–622.
- [R] S. Rueda, Some algebras similar to the enveloping algebra of $\mathfrak{sl}_2$, *Comm. Algebra* **30** (2002), 1127–1152.
- [S] S. P. Smith, A class of algebras similar to the enveloping algebra of $\mathfrak{sl}_2$, *Trans. Amer. Math. Soc* **322** (1990), 285–314.

Department of Mathematics
Franklin and Marshall College, Lancaster, PA 17604
iwan.praton@fandm.edu